

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
19 April 2001 (19.04.2001)

PCT

(10) International Publication Number  
**WO 01/27322 A2**

- (51) International Patent Classification<sup>7</sup>: C12Q 1/68 (81) Designated States (*national*): AE, AG, AL, AM, AT, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KR (utility model), KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (21) International Application Number: PCT/US00/21048
- (22) International Filing Date: 2 August 2000 (02.08.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/159,221 13 October 1999 (13.10.1999) US (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): ELITRA PHARMACEUTICALS, INC. [US/US]; Suite A, 3510 Dunhill Street, San Diego, CA 92121 (US).
- (72) Inventor; and
- (75) Inventor/Applicant (*for US only*): ZYSKIND, Judith [US/US]; 8514 La Jolla Scenic Drive, La Jolla, CA 92047 (US).
- (74) Agent: HUNT, Dale, C.; 16th Floor, 620 Newport Center Drive, Newport Beach, CA 92660 (US).
- Published:  
*Without international search report and to be republished upon receipt of that report.*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: CHITOBIASE AS A REPORTER ENZYME

(57) Abstract: The present invention relates to reporter gene constructs encoding a cytoplasmic form of chitobiase (N,N'-diacetylchitobiase) and methods of using these reporter gene constructs. The use of a cytoplasmic form of chitobiase as a reporter enzyme is generally applicable to the study of gene expression in organisms which do not contain N-acetyl- $\beta$ -D-glucosaminidases.

WO 01/27322 A2

**CHITOBIASE AS A REPORTER ENZYME****GOVERNMENT INTEREST IN THE INVENTION**

Work performed in developing the present invention was supported, in part, by National Science Foundation  
5 grant MCB-9507209, NIH MBRS grant GM45765, and NIH NIGMS MARC F31 GM14967-0451. Accordingly, the  
United States Government may have certain rights in the invention.

Background of the InventionField of the Invention

Reporter enzymes are enzymes whose activities are easily assayed when present inside cells. In order to  
10 study the regulation of a gene whose expression is regulated by various environmental and/or cellular factors or  
influences, a gene encoding a reporter enzyme may be fused to the coding region or to the regulatory region of the  
regulated gene.

One of the most popular cytoplasmic reporter enzymes for use in bacteria is  $\beta$ -galactosidase. It is widely  
used in the art; however, because bacteria such as *Escherichia coli* contain an endogenous  $\beta$ -galactosidase encoding  
15 gene and  $\beta$ -galactosidase may be present in the cytoplasm of such bacteria, deletions of the LacZ gene, the source of  
the enzyme, must be introduced into the host cell line prior to its use. One goal of the present invention is to provide  
an alternative intracellular enzyme for use as a reporter.

This invention relates to genetic constructs and methods of using a cytoplasmic form of the chitobiase  
enzyme as a reporter. The invention also comprises expression vectors which express the cytoplasmic form of  
20 chitobiase. As used herein, all instances of the terminology "chitobiase" refer to a form of chitobiase which is present  
in the cytoplasm of the cell. Cytoplasmic forms of chitobiase may be generated via genetic engineering or microbial  
selection techniques.

Summary of the Invention

One embodiment of the present invention is a method for characterizing a promoter comprising providing a  
25 construct comprising the promoter operably linked to a nucleic acid encoding a cytoplasmic form of chitobiase,  
introducing the construct into host cells, and identifying sequences in the promoter which regulate transcription levels.  
In one aspect of this embodiment, the cytoplasmic form of chitobiase lacks a signal sequence. In another aspect of  
this embodiment, the nucleic acid encoding a cytoplasmic form of chitobiase encodes a fusion protein which comprises  
a cytoplasmic form of chitobiase fused to a heterologous polypeptide. In still another aspect of this embodiment, the  
30 nucleic acid encoding a cytoplasmic form encodes a cytoplasmic form of chitobiase obtained from an organism  
selected from the group consisting of *Alteromonas sp. O-7*, *Arabidopsis thaliana*, *Bacillus subtilis*, *Bombyx mori*, *Bos*  
*taurus*, *Caenorhabditis elegans*, *Candida albicans*, *Dictyostelium discoideum*, *Entamoeba histolytica*, *Felis catus*, *Homo*  
*sapiens*, *Korat cats*, *Lactobacillus casei*, *Leishmania donovani*, *Mus musculus*, *Pisum sativum*, *Porphyromonas*  
*gingivalis*, *Pseudoalteromonas sp. S9*, *Rattus norvegicus*, *Serratia marcescens*, *Streptomyces plicatus*, *Streptomyces*  
35 *thermofilaceus*, *Sus scrofa*, *Trichoderma harzianum*, *Vibrio furnissii*, *Vibrio harveyi*, *Vibrio parahaemolyticus*, and

*Vibrio vulnificus*. In a further aspect of this embodiment, the method of identifying sequences which are involved in directing transcription comprises mutagenizing the promoter. In another aspect of this embodiment, the method of identifying sequences which are involved in transcription comprises constructing deletions in the promoter.

Another embodiment of the present invention is a method for identifying a regulatory element capable of directing or regulating transcription within a test nucleic acid sequence comprising providing a construct comprising the test nucleic acid sequence operably linked to a nucleic acid encoding a cytoplasmic form of chitinase, introducing the construct into host cells, and determining the level of chitinase activity. In one aspect of this embodiment, the cytoplasmic form of chitinase lacks a signal sequence. In another aspect of this embodiment, the nucleic acid encoding a cytoplasmic form of chitinase encodes a fusion protein, the fusion protein comprising a cytoplasmic form of chitinase fused to a heterologous polypeptide. In still another aspect of this embodiment, the nucleic acid encoding a cytoplasmic form encodes a cytoplasmic form of chitinase obtained from an organism selected from the group consisting of *Alteromonas* sp. O-7, *Arabidopsis thaliana*, *Bacillus subtilis*, *Bombyx mori*, *Bos taurus*, *Caenorhabditis elegans*, *Candida albicans*, *Dictyostelium discoideum*, *Entamoeba histolytica*, *Felis catus*, *Homo sapiens*, *Korat cats*, *Lactobacillus casei*, *Leishmania donovani*, *Mus musculus*, *Pisum sativum*, *Porphyromonas gingivalis*, *Pseudoalteromonas* sp. S9, *Rattus norvegicus*, *Serratia marcescens*, *Streptomyces plicatus*, *Streptomyces thermoviolaceus*, *Sus scrofa*, *Trichoderma harzianum*, *Vibrio furnissii*, *Vibrio harveyi*, *Vibrio parahaemolyticus*, and *Vibrio vulnificus*. In one aspect of this embodiment, the reporter gene construct is introduced transiently. In another aspect of this embodiment, the reporter gene construct is introduced stably. The host cells may be selected from the group consisting of prokaryotic cells and eukaryotic cells. In another aspect of this embodiment, the method further comprises permeabilizing or lysing the host cells. The permeabilizing or lysing step may comprise treating the host cells with toluene. The step of determining the level of chitinase activity may be selected from the group consisting of measuring the amount of a chemiluminescent product produced from a substrate, measuring the amount of a fluorescent product produced from a substrate, measuring the amount of light absorbed by a product produced from a substrate and measuring a decrease in the amount of a detectable substrate. In another embodiment, the step of determining the level of chitinase activity may comprise determining the level of *p*-nitrophenol released from a substrate. In another aspect of this embodiment, the test nucleic acid sequence comprises a portion of genomic DNA. In a further aspect of this embodiment, the step of determining the level of chitinase activity comprises determining the level of chitinase activity after exposing the host cells to a desired set of environmental conditions. In still another aspect of this embodiment, the step of determining the level of chitinase activity comprises determining the level of chitinase activity after contacting the host cells with a compound to be tested for its influence on the level of transcription from said regulatory element.

Another embodiment of the present invention is a method of detecting successful transformation, comprising the steps of introducing a nucleic acid encoding a cytoplasmic form of chitinase into host cells, and detecting chitinase expression in the host cells.

Another embodiment of the present invention is a fusion protein-reporter gene construct comprising a promoter operably linked to a nucleic acid encoding a cytoplasmic form of chitinase fused in frame with a nucleic acid encoding a heterologous polypeptide, wherein the heterologous polypeptide is not  $\beta$ -galactosidase or a portion thereof, and wherein the heterologous polypeptide does not contain a signal peptide. In one aspect of this embodiment, the nucleic acid encodes a cytoplasmic form of chitinase obtained from an organism selected from the group consisting of *Alteromonas* sp. O-7, *Arabidopsis thaliana*, *Bacillus subtilis*, *Bombyx mori*, *Bos taurus*, *Caenorhabditis elegans*, *Candida albicans*, *Dictyostelium discoideum*, *Entamoeba histolytica*, *Felis catus*, *Homo sapiens*, *Korat cats*, *Lactobacillus casei*, *Leishmania donovani*, *Mus musculus*, *Pisum sativum*, *Porphyromonas gingivalis*, *Pseudoalteromonas* sp. S9, *Rattus norvegicus*, *Serratia marcescens*, *Streptomyces plicatus*, *Streptomyces thermoviolaceus*, *Sus scrofa*, *Trichoderma harzianum*, *Vibrio furnissii*, *Vibrio harveyi*, *Vibrio parahaemolyticus*, and *Vibrio vulnificus*. In another aspect of this embodiment, the nucleic acid further comprises a  $\lambda$  site-specific recombination sequence.

Another embodiment of the present invention is a reporter gene construct comprising plasmid pJMF3.

Another embodiment of the present invention is a reporter gene construct comprising plasmid pJMF4.

15 Another embodiment of the present invention is a reporter gene construct comprising plasmid pDYK9.

Another embodiment of the present invention is a reporter gene construct comprising plasmid pDYK11.

Another embodiment of the present invention is a host cell comprising a fusion protein-reporter gene construct comprising a promoter operably linked to a nucleic acid encoding a cytoplasmic form of chitinase fused in frame with a nucleic acid encoding a heterologous polypeptide, wherein the heterologous polypeptide is not  $\beta$ -galactosidase or a portion thereof, and wherein the heterologous polypeptide does not contain a signal peptide. In one aspect of this embodiment, the nucleic acid is integrated into a chromosome of the cell. In another aspect of this embodiment, the nucleic acid is transiently expressed in the host cell.

Another embodiment of the present invention is a nucleic acid encoding a cytoplasmic form of chitinase in which the signal sequence of native chitinase has been inactivated or deleted. In one aspect of this embodiment, the signal sequence has been mutated to inactivate it.

An isolated or purified polypeptide comprising a cytoplasmic form of chitinase fused in frame with a heterologous polypeptide, wherein the heterologous polypeptide is not  $\beta$ -galactosidase or a portion thereof and wherein the heterologous polypeptide does not contain a signal peptide.

Another embodiment of the present invention is an isolated or purified polypeptide comprising a cytoplasmic form of chitinase in which the signal peptide of native chitinase has been inactivated or deleted. In one aspect of this embodiment, the signal sequence has been mutated to inactivate it.

Another embodiment of the present invention is a method for monitoring the activity of a promoter comprising providing a construct comprising the promoter operably linked to a nucleic acid encoding a cytoplasmic form of chitinase, introducing the construct into host cells, and determining the level of chitinase activity. In one aspect of this embodiment, the cytoplasmic form of chitinase lacks a signal sequence. In another aspect of this

embodiment, the nucleic acid encoding a cytoplasmic form of chitobiase encodes a fusion protein, the fusion protein comprising a cytoplasmic form of chitobiase fused to a heterologous polypeptide. In one aspect of this embodiment, the nucleic acid encoding a cytoplasmic form encodes a cytoplasmic form of chitobiase obtained from an organism selected from the group consisting of *Alteromonas sp. O-7*, *Arabidopsis thaliana*, *Bacillus subtilis*, *Bombyx mori*, *Bos taurus*, *Caenorhabditis elegans*, *Candida albicans*, *Dictyostelium discoideum*, *Entamoeba histolytica*, *Felis catus*, *Homo sapiens*, *Korat cats*, *Lactobacillus casei*, *Leishmania donovani*, *Mus musculus*, *Pisum sativum*, *Porphyromonas gingivalis*, *Pseudoalteromonas sp. S9*, *Rattus norvegicus*, *Serratia marcescens*, *Streptomyces plicatus*, *Streptomyces thermoviolaceus*, *Sus scrofa*, *Trichoderma harzianum*, *Vibrio furnissii*, *Vibrio harveyi*, *Vibrio parahaemolyticus*, and *Vibrio vulnificus*. In another aspect of this embodiment, the reporter gene construct is introduced transiently. In still another aspect of this embodiment, the reporter gene construct is introduced stably. In a further aspect of this embodiment, the host cells are selected from the group consisting of prokaryotic cells and eukaryotic cells. In yet another aspect of this embodiment, the method further comprises permeabilizing or lysing the host cells. For example, the permeabilizing or lysing step may comprise treating the host cells with toluene. In yet another aspect of this embodiment, the step of determining the level of chitobiase activity may be selected from the group consisting of measuring the amount of a chemiluminescent product produced from a substrate, measuring the amount of a fluorescent product produced from a substrate, measuring the amount of light absorbed by a product produced from a substrate and measuring a decrease in the amount of a detectable substrate. In still another aspect of this embodiment, the step of determining the level of chitobiase activity comprises determining the level of *p*-nitrophenol released from a substrate. In still another aspect of this embodiment, the step of determining the level of chitobiase activity comprises determining the level of chitobiase activity after exposing the host cells to a desired set of environmental conditions. In a further aspect of this embodiment, the step of determining the level of chitobiase activity comprises determining the level of chitobiase activity after contacting the host cells with a compound to be tested for its influence on the level of transcription from said regulatory element. For example, the compound may comprise a compound to be tested for activity as a drug.

#### 25 Brief Description of the Drawings

Figure 1 illustrates plasmids pJMF3 and pJMF4 containing *attP* in 2 different orientations and the *lac* promoter with the first 21 amino acids of *lacZα* (from pUC19) fused in-frame to the *chb* gene. The sequence of the fusion region is shown in Figure 2. Restriction sites shown are found once in the plasmid sequences except for *NotI*, which has 2 sites flanking the P15A origin.

30 Figure 2 illustrates the sequence of the *lac* promoter and the chitobiase fusion found in pJMF3 and pJMF4 (SEQ ID NOS: 15 and 16). Fusion between *lacZα* (from pUC19) and *chb* [Soto-Gill, R.W. et al, J. Biol. Chem. 264:14778-14782 (1998)] is indicated by (/); start of transcription is indicated by (+1). Sequence and binding sites in the *lac* promoter regulatory region are found in Dickson, R.C. et al., Science 187:27-32 (1975). Restriction enzyme sites shown in Figure 2 are found once in the plasmid sequences except for *SphI*, which has 2 sites; these different

restriction sites can be used to replace the *lac* promoter with another promoter together with part of a coding region to produce an in-frame fusion with *chb*.

Figure 3 depicts integration of *chb* fusions into the chromosome by site-specific recombination between *attB* and *attP*. The steps involved are described below and in Diederich, L.L. et al., Plasmid 28:14-24 (1992).

5        Figure 4 illustrates plasmids pDYK9 and pDYK11 which contain *dnaA-chb* fusions. pDYK9 is deleted for the *rpmH* regulatory region. The orientation of *attP* in pDYK9 and pDYK11 is the same as that of pJMF3. After integration at *attB* of the larger *NotI* fragment, the *dnaA* promoters are oriented to transcribe in the same direction as replication fork movement.

10        Figure 5 illustrates the *rpmH-dnaA* regulatory region and *dnaA*-chitobiase fusion (SEQ ID NO.: 17). The *dnaA* box and promoters are shaded, and the coding region of the *dnaA* gene is striped. The fusion contains two amino acids (between the backslashes) from pUC19. The region cloned into pDYK9 is between *SphI* primer II and *KpnI* primer, and the region cloned into pDYK11 is between *SphI* primer I and *KpnI* primer. The numbers above the primers refer to the nucleotide in the sequence amplified. For sequence numbering and the locations of promoters and protein binding sites, see Froelich, J.M. et al., J. Bacteriol. 178:6006-6012 (1996); Hansen, F.G. et al., EMBO J. 1:1043-1048 (1982) and Messer, W. and C.W. Weigel, Initiation of Chromosome Replication. p. 1579-1601 (1996) In F.C. Neidhart, R. Curtiss III, J.L. Ingraham, E.C.C. Lin, K.B. Low, B. Magasanik, W.S. Reznikoff, M. Riley, M. Schaechter and H.E. Umbarger (Eds.), Escherichia coli and Salmonella Cellular and Molecular Biology., ASM Press, Washington, D.C.

#### Detailed Description of the Preferred Embodiment

20        Reporter genes and reporter gene constructs play a number of important roles in a variety of molecular biology techniques. For example, reporter genes may be used to determine whether a sequence contains a promoter or other cis-acting element which directs transcription, such as an enhancer. In addition, reporter genes may be used to identify regulatory sites in promoters or other cis-acting elements and to determine the effects of mutating these regulatory sites on the level of gene expression directed by the promoters or other cis-acting elements. Reporter genes may also be used to detect successful transformation. In addition, reporter genes may be used to monitor gene expression under various conditions and to identify drugs.

30        The structure of a reporter gene construct containing the cytoplasmic form of chitobiase will vary according to its purposes. When the reporter gene construct is a vector, one must decide between a vector that incorporates the reporter gene into the host's genome or one that replicates extrachromosomally, such as a plasmid. When the reporter construct is not designed to integrate into the host genome, the vector can contain an origin of replication with activity in the host cell of interest. This feature provides the reporter gene vector the ability to replicate within the host cell in which it has been introduced. In addition to the origin of replication, reporter constructs often contain a promoter, a multiple cloning site, a selectable marker, and of course a reporter gene. Reporter constructs for use in eukaryotic cells may also contain a polyA site adjacent to the reporter gene.

Given the utility of reporter gene constructs, it is not surprising that a number of cytoplasmic reporter gene constructs and different reporter genes are available for use by those of skill in the art. For example, the cytoplasmic reporter enzymes chloramphenicol acetyltransferase (CAT), firefly luciferase,  $\beta$ -glucuronidase (GUS), green fluorescent protein (GFP), and  $\beta$ -galactosidase have been used extensively. However, such reporters all have individual shortcomings that may limit or preclude their usage under some conditions. For example, high levels of GFP are toxic to the cell. In addition, reporter enzymes are not expressed equally in all cell types nor are they equally stable when expressed in all cell types. Furthermore, there is a recognized need for multiple reporter enzymes that can be assayed independently of one another in order to simultaneously study the regulation of multiple genes within a single cell type. Therefore, there exists a continuing need to identify reporter enzymes with useful properties.

The cytoplasmic enzyme  $\beta$ -galactosidase is widely used as a reporter gene in various microbiological and molecular biological studies. This enzyme is used in both *in vitro* and *in vivo* assays. The wide acceptance of this reporter system results, in part, because it is non-isotopic and extremely flexible. It is used in a number of assay formats and has an extremely broad linear range. Nevertheless, because  $\beta$ -galactosidase is present in the cytoplasm of various host cells such as *Escherichia coli*, deletion of the *lacZ* gene, the source of the enzyme, is often required prior to its use in a host cell system. One goal of the present invention was to provide an alternative intracellular enzyme for use as a reporter.

An extensive discussion of various molecular biology techniques is available in Ausubel, et al., (eds) "Short Protocols in Molecular Biology," Wiley and Sons, Inc., New York (1997). Examples of such techniques include isolating and preparing DNA for manipulation, gel electrophoresis, polymerase chain reaction (PCR), determining nucleic acid sequences, screening nucleic acid libraries, mutagenesis of DNA, and introducing DNA into host cells.

The structure of a reporter gene construct will vary according to its purposes. The reporter gene constructs are constructed according to standard techniques of molecular biology well known in the art.

When the reporter gene construct is a vector, one must decide between a vector that incorporates the reporter gene into the host's genome or one that replicates extrachromosomally, such as a plasmid. When integration of the reporter gene is a desired result, a reporter gene construct will contain sequences that will facilitate incorporation.

One example of integration sequences that can be included in a reporter gene construct is the  $\lambda$  *attP* site. This site permits a single copy of the reporter gene to be incorporated into a host bacterial genome. Integration-promoting sequences with utility in mammalian cells include the long terminal repeats found in retroviral genomes. These sequences promote viral genome integration in a host genome and have been used extensively by those of skill in the art to promote the integration of exogenous sequences in mammalian host cells.

When the reporter construct is not designed to integrate into the host genome, it is common that the vector contain an origin of replication with activity in the host cell of interest. This feature provides the reporter gene vector the ability to replicate within the host cell in which it has been introduced.

In addition to the origin of replication, reporter constructs will often contain additional features that promote the expression of the nucleic acid sequence or sequences contained in the construct. These additional sequences can include a polyA site, a multiple cloning site, a drug resistance marker, and of course a reporter gene.

The present invention relates to the use of chitinase as a reporter gene. The chitinase may be used as a reporter in bacteria, plants, mammalian cells and other host cell lines. One possible alternative to using  $\beta$ -galactosidase as a reporter gene was to develop cytoplasmic  $N,N'$ -diacetylchitinase ( $N$ -acetyl- $\beta$ -D-glucosaminidase, EC 3.2.1.30) for use as a reporter enzyme. One advantage of the enzyme  $N$ -acetyl- $\beta$ -D-glucosaminidase or "chitinase" over  $\beta$ -galactosidase is that genes encoding chitinase are missing from many bacteria, including *E. coli*, some fungi, and some eukaryotic cells. Thus, it is not necessary to engineer many host cells to lack reporter activity as is the case with  $\beta$ -galactosidase. The present invention also relates to various protein expression vectors that can be used to express the reporter gene. In addition, the present invention may be used in conjunction with other reporter enzymes in systems in which the regulation or activities of multiple genes is to be studied simultaneously.

Chitinase is one of two enzymes that hydrolyze chitin, an abundant insoluble polysaccharide, to its monomeric unit,  $N$ -acetylglucosamine (GlcNAc). Chitinase is known to be present in a number of organisms. For example, the chitinase enzyme is known to be present in various genera including *Arabidopsis*, *Bacillus*, *Bombyx*, *Bos*, *Caenorhabditis*, *Candida*, *Dictyostelium*, *Entamoeba*, *Felis*, *Homo*, *Korat*, *Lactobacillus*, *Leishmania*, *Mus*, *Pisum*, *Porphyromonas*, *Pseudoalteromonas*, *Rattus*, *Serratia*, *Streptomyces*, *Sus*, *Trichoderma*, and *Vibrio*. Specific examples of organisms known to contain chitinase include *Alteromonas* sp. O-7, *Arabidopsis thaliana*, *Bacillus subtilis*, *Bombyx mori*, *Bos taurus*, *Caenorhabditis elegans*, *Candida albicans*, *Dictyostelium discoideum*, *Entamoeba histolytica*, *Felis catus*, *Homo sapiens*, *Korat cats*, *Lactobacillus casei*, *Leishmania donovani*, *Mus musculus*, *Pisum sativum*, *Porphyromonas gingivalis*, *Pseudoalteromonas* sp. S9, *Rattus norvegicus*, *Serratia marcescens*, *Streptomyces plicatus*, *Streptomyces thermoviolaceus*, *Sus scrofa*, *Trichoderma harzianum*, *Vibrio furnissii*, *Vibrio harveyi*, *Vibrio parahaemolyticus*, and *Vibrio vulnificus*.

One source of the enzyme is the marine bacterium, *Vibrio harveyi*. *Escherichia coli* cells harboring a plasmid carrying the *chb* gene from *Vibrio harveyi* were reported to produce the enzyme, which was found to be associated with the outer membrane of the bacterial cells. These are described in Jannatipour, M. et al., "Translocation of *Vibrio harveyi*  $N,N'$ -diacetylchitinase to the outer membrane of *Escherichia coli*," J. Bacteriol. 169:3785-3791 (1987) and Soto-Gil & Zyskind,  $N,N'$ -Diacetylchitinase of *Vibrio harveyi* primary structure, processing, and evolutionary relationships. J. Biol Chem. 264:14778-14782 (1989).

The present invention contemplates expressed cytoplasmic forms of chitinase in various forms. In one embodiment, the signal sequence is deleted from the amino terminal portion of the protein. Presumably the removal of this sequence results in the expression of a cytoplasmic form of the enzyme that is not secreted from the host cell or incorporated into the membrane of the host cell producing the enzyme.

The present invention also contemplates the generation of fusion proteins comprising a fusion polypeptide joined in frame to chitinase. Preferably, the fusion polypeptide comprises a polypeptide other than chitinase, such



as a heterologous protein. The heterologous polypeptide may comprise a polypeptide having a biological activity (such as an enzymatic or other activity besides activity as an immunogen) or the heterologous polypeptide may not have a biological activity. The heterologous polypeptide does not include a signal sequence which directs its secretion. Preferably the heterologous polypeptide is not  $\beta$ -galactosidase or a portion thereof. Thus, the fusion reporter gene construct contains a sequence encoding the fusion polypeptide genetically fused in frame with a sequence encoding chitobiase. In one embodiment, this fusion may remove the amino-terminal signal peptide sequence of chitobiase and replace it with a heterologous protein.

In another embodiment, the fusion protein construct comprises a chitobiase gene sequence that has been truncated to remove at least the signal peptide sequence of the gene. Alternatively, mutations may be introduced into the signal peptide sequence so that it is no longer functional. Such mutations may be introduced using a variety of techniques familiar to those skilled in the art, including site directed mutagenesis, cassette mutagenesis, and chemical mutagenesis.

Once the reporter gene construct is made it is introduced into a host cell line for testing. Host cells of prokaryotic and eukaryotic origin can be used with the reporter gene constructs of the present invention. A variety of methods are available to introduce reporter gene constructs into host prokaryotic cells. For example, bacteria can be transformed using calcium chloride, electroporation, or viral vectors such as the filamentous phages. These and other prokaryotic transformation protocols are well known in the art.

Alternatively, the sequence encoding chitobiase may be introduced in eukaryotic cells, including yeast, mammalian, plant, and insect cells. For example, the sequence encoding chitobiase may be inserted into a yeast artificial chromosome, a yeast plasmid, a bovine papilloma virus vector or other extrachromosomal vector, a retroviral vector, a Ti-plasmid, or a baculovirus vector. A variety of such vectors are known to those skilled in the art. The vectors may be introduced into any of the yeast, mammalian, plant, and insect cells familiar to those skilled in the art.

The introduction of the reporter gene construct into mammalian cells can likewise utilize a number of transfection protocols well known to those of skill in the art. As discussed above, transfections can be transient or stable. Examples of suitable transfer protocols include calcium phosphate transfection, DEAE-Dextran, electroporation, liposome-mediated transfection, and viral transfection. These and other eukaryotic transformation protocols are well known in the art.

Following introduction of the reporter gene construct into the host cell of interest, the enzymatic activity of the reporter gene is measured. Preferably, the chitobiase assays are performed after permeabilizing or lysing the host cells. There are a variety of cell permeabilization and cell lysis procedures available to those of ordinary skill in the art, including methods such as sonication or lysozyme treatment. One embodiment of the present invention uses toluene treatment to permeabilize cells. The details of this method are discussed in D.Y. Kalabat et al., BioTechniques 25:1030-1035 (1998) and Miller, J.H. A Short Course in Bacterial Genetics, CSH Laboratory Press, Cold Spring Harbor, NY 1992.

Cellular chitinase activity can be measured quantitatively by following the hydrolysis of chitinase substrates. Examples of substrates with utility in chitinase activity assays include *N,N'*-diacetylchitinase (chitinase), *p*-nitrophenyl-*N*-acetyl- $\beta$ -D-glucosaminide (PNAG)(Sigma Chemical, St. Louis, MO), and 5-bromo-4-chloro-3-indolyl-*N*-acetyl- $\beta$ -D-glucosaminide (X-Gluc)(Sigma Chemical, St. Louis, MO). Other substrates are also contemplated for use in the assays of the present invention.

Products produced by the hydrolysis of the chitinase substrates are monitored using various means familiar to those skilled in the art. For example, various optical means are known to those skilled in the art. One such optical means may comprise detection of chemiluminescent or fluorescent products released from a substrate. Alternatively, the level of chitinase activity may be determined by measuring the amount of light absorbed by a product produced from a substrate or measuring a decrease in the amount of a detectable substrate. In one embodiment, *p*-nitrophenol is released from the substrate and measured at 400 nm. Other monitoring methods well known in the art can be used to quantitate signals produced in the chitinase assay. These may include use of radioactive substrates or substrates having radiofrequency tags. In another embodiment, blue/white colony indicator plates are used to monitor enzyme activity.

Another embodiment of the present invention is a kit. One aspect of this embodiment includes a reporter gene construct comprising a vector containing a chitinase reporter gene. The reporter gene construct also contains a multiple cloning site containing a variety of restriction endonuclease cutting sites that facilitate the introduction of exogenous DNA into the construct.

The kit embodiment of the present invention also includes those components necessary to assay for chitinase activity produced by the reporter gene construct. For example, in one embodiment, the kit will include a supply of a suitable chitinase substrate whose metabolism into product by the reporter enzyme can be assayed.

#### EXAMPLES

The following Examples are disclosed to assist in the understanding of the present invention. The Examples below should not be construed to limit the scope of the invention and such variations of the invention now known or later developed, which would be within the purview of one of ordinary skill in the art and are considered to fall within the scope of the invention hereinafter claimed.

##### Example 1

##### Construction of Vectors

Vectors were constructed using polymerase chain reaction (PCR) products that were cloned first at the *Sma*I or *Eco*RV restriction sites of plasmid pBluescript II (pKSII+)(Stratagene; San Diego, CA). The nucleotide sequence of all PCR products was determined using standard techniques well known in the art. Unmethylated plasmid DNA was isolated from an *E. coli dam*<sup>-</sup> strain when cutting with the *Bcl*I enzyme was required.

The pDYK9 (SEQ ID NO: 11) plasmid was constructed by ligating a *Sph*I-*Kpn*I PCR product containing the *dnaA* promoter region into plasmid pRSG196 that contains the *V. harveyi* chitinase gene (see Jannatipour, M., et al.). Briefly, the *V. harveyi* chitinase gene was prepared by cloning a *Sau*3A partial digest of *V. harveyi* DNA into the

single *Bam*HI site with the *tet* gene of pMK2004 [See Soto-Gill & Zyskind, "Cloning of *Vibrio harveyi* chitinase and chitobiase genes in *Escherichia coli*," pp. 209-223, in J.P. Zikakis (ed.), Chitin, chitosan, and related enzymes," Academic Press, Inc., New York (1984)]. The clones containing the gene of interest were detected by the presence of the yellow *p*-nitrophenylate product after individual colonies of the clone bank were sprayed with 10 mM PNAG dissolved in 100 mM sodium phosphate, pH 7.0. Restriction maps of the positive clones were made according to standard techniques well known in the art.

The *chb* gene was subcloned in a 3.5-kb *Eco*RI fragment from pRSG14 [See Jannatipour, M. et al., "Translocation of *Vibrio harveyi* *N,N'*-diacetylchitobiase to the outer membrane of *Escherichia coli*," J. Bacteriol. 169:3785-3791 (1987)] into the *Eco*RI site of pUC19 [Yanisch-Perron, c.J. et al., Gene 33: 103-119 (1985)]. Plasmid pRSG196 was constructed by deleting a 0.5-kb *Sst*I fragment from one of these clones [See Jannatipour, M. et al., "Translocation of *Vibrio harveyi* *N,N'*-diacetylchitobiase to the outer membrane of *Escherichia coli*," J. Bacteriol. 169:3785-3791 (1987)].

The pRSG196 plasmid was also cut with *Sph*I and *Kpn*I to accommodate the insertion of the *dnaA* fragment. Plasmid pAC17 [described in Chiaramello & Zyskind, "Coupling of DNA replication to growth rate in *Escherichia coli*: a possible role for guanosine tetraphosphate," J. Bacteriol. 172:2013-2019 (1992)] served as template with primers 5'-GCA CAT GCA TGC TGG TCA TTA AAT TTT CC-3' (SEQ ID NO 1) and 5'-CGG GGT ACC AAC TCA TCC TGC AAT CG-3' (SEQ ID NO 2) producing a PCR product 374 bp long that contains 353 bp from the *dnaA* promoter region [bases 583 to 935, numbering according to Hansen, F. G., et al., "The nucleotide sequence of the *dnaA* gene promoter and of the adjacent *rpmH* gene, coding for the ribosomal protein L34, of *Escherichia coli*," EMBO J. 1:1043-1048 (1982)]. The forward primer (*Sph*I primer II) contained an *Sph*I site and the reverse primer (*Kpn*I primer) contained a *Kpn*I site for cloning. This created an in-frame fusion between the amino-terminal 17 amino acids of DnaA and the carboxy-terminal end of chitobiase deleted for the amino-terminal 22 amino acids including the signal peptide.

The next step in constructing the pDYK9 vector was taking the 3270 bp *Dra*I-*Hind*III (partial digest) fragment containing the *dnaA-chb* fusion and ligating it to the pACYC184 2555 bp *Hinc*II-*Hind*III fragment carrying chloramphenicol-resistance (*Cm<sup>r</sup>*) and the P15A origin. This fragment was described in Chang & Cohen, "Construction and characterization of amplifiable multicopy DNA cloning vehicles derived from the p15A cryptic miniplasmid," J. Bacteriol. 134:1141-1156 (1987).

A *Not*I site was introduced at the *Acc*I site after digestion with *Acc*I, treatment with Mung bean nuclease, and ligation to phosphorylated *Not*I linkers (New England Biolabs, Inc.; Beverly, MA). A *Not*I site was introduced at the *As*eI site after digestion with *As*eI, treatment with Mung bean nuclease, and ligation to phosphorylated *Not*I linkers (New England Biolabs, Inc.; Beverly, MA). An *Xba*I-*Sph*I PCR product containing the *rrnBt1t2* terminator was ligated into this plasmid cut with the same enzymes creating pDYK7. *E. coli* chromosomal DNA served as template with primers 5'-CTA GTC TAG ATG CCG AAC TCA GAA GTG A-3' (SEQ ID NO 3) and 5'-GCA CAT GCA TGC GGG GGA TGG CTT GTA GAT-3' (SEQ ID NO 4) to produce a PCR product 357 bp long that contains bases 6534 to 6869 from the *rrnB* operon [numbering according to Brosius, J., et al., "Gene organization and primary structure of a

ribosomal RNA operon from *Escherichia coli*." J. Mol. Biol. 148:107-127 (1981)) and includes the complex transcription termination region of this operon which is described in Orosz, A., et al., "Analysis of the complex transcription termination region of the *Escherichia coli rrmB* gene," Eur. J. Biochem. 201:653-659 (1991). The forward primer contained an *Xba*I site and the reverse primer contained an *Sph*I site.

5 A *Bcl*-*Sma*I PCR product containing the  $\lambda$  *attP* site was ligated into pDYK7 cut with *Tth*111I, treated with Mung bean nuclease, then digested with *Bcl*I. Plasmid pHN894 [described in Goodman, S. D., et al., "Deformation of DNA during site-specific recombination of bacteriophage lambda: Replacement of IHF protein by HU protein or sequence-directed bends," Proc. Natl. Acad. USA 89:11910-11914 (1992)] served as template with the forward primer 5'-CAT GAT CAT GCG ACA GGT TTG ATG A-3' (SEQ ID NO 5) and the reverse primer 5'-GGG GGC GCC TAC  
10 CTT TCA CGA G-3' (SEQ ID NO 6) producing a PCR product 466 bp long that contains the  $\lambda$  *attP* site. The 466 bp PCR product containing the  $\lambda$  *attP* site was first cloned into the *Sma*I site of the pKSII+ plasmid to produce pDYK8. The forward primer contains a *Bcl*I site and the reverse primer contains G's at the 5' end in order to recreate a *Sma*I site when cloning the PCR fragment into a *Sma*I site. This PCR product includes bases -211 to +241 from the center of the *attP* core and the sequence required for optimum  $\lambda$  *attP* site integration as described in Nagaraja & Weisberg,  
15 "Specificity determinants in the attachment sites of bacteriophages HK022 and  $\lambda$ ," J. Bacteriol. 172:6540-6550 (1990). The orientation of *attP* is such that when the fusion is integrated at *attB*, the transcription direction of *dnaA**p1* and *p2* promoters is the same as replication fork movement mimicking the orientation at the wild type *dnaA* promoters.

Vector pDYK11 (SEQ ID NO: 12) was constructed by ligating an *Sph*I-*Kpn*I PCR product containing the *rpmH*-  
20 *dnaA* promoter region into pDYK9 also digested with *Sph*I and *Kpn*I. Plasmid pAC17, described in Chiamarello & Zyskind, (1992), served as template with the primers 5'-CAT GCA TGC ATG AAA CGA TGG ACA CC-3' (SEQ ID NO 7) and 5'-CGG GGT ACC AAC TCA TCC TGC AAT CG-3' (SEQ ID NO 8) to produce a PCR product 616 bp long that contains 598 bp from the *rpmH*-*dnaA* regulatory region [bases 338 to 935, numbering according to Hansen, F. G., et al., (1982)]. The forward primer (*Sph*I primer I) contained an *Sph*I site and the reverse primer (*Kpn*I primer) contained a  
25 *Kpn*I site for cloning. SEQ ID NO: 18 provides the complete coding sequence of the DNA encoding the *dnaA*/chitobiase fusion protein. The complete sequence of the *dnaA*/chitobiase fusion protein is provided in SEQ ID NO: 19.

The vector pJMF3 (SEQ ID NO: 13) was constructed by first ligating an *As*eI linker (5'-CATTAATGCATG-3' (SEQ ID NO 9) self-hybridized) into the *Sph*I site of pDYK11. The pUC19, [described in Yanisch-Perron C., et al., "Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors,"  
30 Gene 33:103-19 (1985)] *As*eI-*Kpn*I fragment containing the *lacPO*-polylinker region was ligated to this plasmid after digestion with *As*eI and *Kpn*I. The resulting in-frame fusion between the amino-terminal 21 amino acids of the pUC19 *lacZ*( $\alpha$ ) peptide [Yanisch-Perron C., et al., 1985] and the carboxy-terminal end of chitobiase deleted for the amino-terminal 22 amino acids is identical to the protein fusion in pRSG196 [Jannatipour, M., et al., (1995)].

Vector pJMF4 (SEQ ID NO: 14 ) was constructed by ligating the *Bam*HI-*Eco*RV fragment of pDYK8  
35 containing the  $\lambda$  *attP* site into pDYK7 which had been digested with *Tth*111I, treated with Mung bean nuclease, then

digested with *Bcl*I, to create pTKP9. This reversed the orientation of the *attP* site relative to the *attP* site in pDYK9. The pDYK11 *Bsp*MI-*Kpn*I fragment containing the *rpmH-dnaA* promoter region was then ligated into pTKP9 digested with the same enzymes to create pTKP11. This reversed the orientation of the *attP* site relative to the *attP* site in pDYK11. An *Asel* linker (5'-CATTAAATGCATG-3' (SEQ ID NO 10) self-hybridized) was ligated into the *Sph*I site of pTKP11 to create pJMF2. The pUC19 [as described in Yanisch-Perron C., et al., (1985)] *Asel-Kpn*I fragment containing the *lacPO*-polylinker region was ligated to pJMF2 cut with *Asel* and *Kpn*I to create pJMF4.

#### Example 2

##### Site-specific Recombination

To move the chitobiase fusions in pDYK9 and pDYK11 to the *attB* site in the chromosome, *Not*I fragments from these plasmids were self-ligated and transformed or electroporated into strain WM2269 (DH5 $\alpha$  containing pLDR8) [See Zyskind, J. W. and S. I. Bernstein (1992) "Recombinant DNA Laboratory Manual," Academic Press, San Diego, CA]. Plasmid pLDR8, described in Diederich, L. L. J., et al., "New cloning vectors for integration into the  $\lambda$  attachment site *attB* of *Escherichia coli* chromosome," Plasmid 28:14-24 (1992), expresses integrase from the  $\lambda$  *P<sub>R</sub>* promoter and contains the  $\lambda$  *cI<sub>857</sub>* repressor gene, a kanamycin resistance gene, and a temperature-sensitive origin of replication. The transformed or electroporated cells were incubated at 42°C with shaking for 30 min then moved to 37°C for 1 h followed by selection on Luria broth agar plates containing chloramphenicol (25  $\mu$ g/ml) at 42°C. Transformants were screened for loss of kanamycin resistance and, therefore, loss of pLDR8.

#### Example 3

##### Bacteriophage P1 Transduction

Transduction with P1 bacteriophage by the method of Zyskind & Bernstein, Recombinant DNA Laboratory Manual. Academic Press, San Diego, CA (1992) was used to construct strains and to confirm the chromosomal location of the *dnaA-chb* fusions. Cotransduction of *Cm<sup>r</sup>* (carried by the fusion) and *galK* (linked to *attB*) indicated that *Cm<sup>r</sup>* and *galK* are linked on the chromosomes of strains DYK9W, DYK9F, DYK11W, and DYK11F.

#### Example 4

##### p-Nitrophenol Chitobiase Activity Assay

Chitobiase activity is located in the cytoplasm when its signal peptide is replaced by fusion with another peptide. Accordingly, chitobiase assays are performed on permeabilized or lysed cells. A variety of permeabilization or cell lysis protocols are available to liberate the enzyme from within a cell population being tested. One such protocol involves toluene-treated cells washed once with M9 salts according to the method of Miller, J.H. A Short Course in Bacterial Genetics, CSH Laboratory Press, Cold Spring Harbor, NY 1992. The toluenized cells are placed in a chitobiase buffer (10 mM Tris-HCl, pH 8.0, and 0.5 M NaCl) and 666  $\mu$ M PNAG, the chitobiase enzyme substrate. NaCl is included because chitobiase has approximately 80% of full activity in the absence of salt, with maximal activity occurring between 0.25 and 0.6 M NaCl. Toluenized cells (0.772 ml) are preincubated at 28°C and the reaction started with the addition of 0.228 ml PNAG (1 mg/ml). After incubation at 28°C, the reaction is stopped by the addition of 1 ml of 1 M Tris base. The release of *p*-nitrophenol is measured at 400 nm and turbidity at 550 nm. *p*-

Nitrophenol release is measured immediately at 400 nm with a molar absorptivity of  $1.8 \times 10^3$  liters  $\text{mol}^{-1} \text{cm}^{-1}$ . Units are calculated after subtracting the light scattering factor ( $1.5 \times \text{OD}_{550}$ ) from  $\text{OD}_{400}$  of the sample. The normalizing factor of 1.5 was determined previously by measuring the light scattering ratio of bacteria at  $\text{OD}_{400}$  and  $\text{OD}_{550}$ . One unit of chitobiase activity is the amount of enzyme that catalyzes the formation of 1 pmol of *p*-nitrophenol per min at 28°C. For comparison to Miller units of  $\beta$ -galactosidase [described in Miller, J. H., A Short Course in Bacterial Genetics. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1992)], the units are normalized to 1 ml of culture at  $\text{OD}_{450} = 1$ .

#### Example 5

##### Determination of Chitobiase Activity in Cells Containing Vectors pJM3 and pJM4

(Figure 1) contain the *lacPO* promoter with the first 21 amino acids of *lacZ* $\alpha$  (from pUC19) fused in-frame to the *chb* gene. These plasmids also contain the  $\lambda$  phage *attP* recombination site in different orientations, the gene encoding chloramphenicol acetyltransferase (*cat*), and a ribosomal terminator, *rrbT1t2*, inserted upstream of the *lac-chb* fusion to prevent read-through from other promoters. The chitobiase activity associated with these plasmids (Table 1) is high in the absence of IPTG because of titration of *lac* repressor expressed from a single copy chromosomal gene. Induction by IPTG is approximately 10-fold (Table 1).

**Table 1. Chitobiase Activity of *lacZ-chb* Fusion<sup>a</sup>**

Plasmid	Chitobiase <sup>b</sup> (U <sup>c</sup> )	
	-IPTG	+IPTG (1 mM)
pJMF3 in DH5 $\alpha$	668 $\pm$ 45	9320 $\pm$ 347
pJMF4 in DH5 $\alpha$	788 $\pm$ 44	7188 $\pm$ 477

<sup>a</sup> Overnight cultures were diluted 1:1000 into 50 mL prewarmed LB and grown to  $\text{OD}_{450} = 0.1$ . 1 mM IPTG was added to half of the culture, and growth continued to  $\text{OD}_{450} = 0.3$ .  
<sup>b</sup> Triplicate samples were assayed. Mean chitobiase activities are given with standard deviations.  
<sup>c</sup> One unit of chitobiase activity is 1 pmol of *p*-nitrophenol/min at 28°C. Units given for 1 mL of culture at  $\text{OD}_{450} = 1$ .

The *lac* promoter can be replaced with another promoter and a fusion protein created with chitobiase by cutting with *SphI* or *AseI* and either *SalI*, *KpnI*, or *SstI* (Figure 2). Fusions created with these vectors can be moved to the chromosome by site specific recombination at the  $\lambda$  *attB* site to permit single copy analysis of the activity of the promoter. The protocol, as described more fully in Diederich, et al., "New cloning vectors for integration into the  $\lambda$  attachment site *attB* of *Escherichia coli* chromosome," Plasmid 28:14-24 (1992), involves two components, (i) a circular DNA containing the  $\lambda$  attachment site, *attP*, the promoter-*chb* gene fusion, and the *cat* gene, and (ii) a helper plasmid, pLDR8, which contains the *int* gene under the control of the temperature-sensitive repressor, *cI857* and a temperature-sensitive origin of replication. The plasmid is digested with *NotI* to remove the P15A origin, and the

fragment containing the chitobiase fusion is self-ligated prior to transformation into cells containing pLDR8. Integration occurs by site-specific recombination between *attP* and *attB* (17.4 min on the *E. coli* chromosome, Figure 3).

#### Example 6

5

#### Use of the Chitobiase Reporter Enzyme to Study *dnaA* Gene regulation

Two plasmids, pDYK9 and pDYK11, discussed in the Examples above, were constructed to assess the regulation of the *dnaA* gene using chitobiase as a reporter enzyme. These plasmids differ by the absence of the *rpmH* regulatory region in pDYK9 (Figure 4). These fusions were moved from the plasmid to the chromosomal *attB* site for single copy analysis as described above. After transformation of strain WM2269 with the ligated DNA, integration occurred by site specific recombination between the *attP* and the *attB* sites. The orientation of *attP* in pDYK9 and pDYK11 is such that when the fusion is integrated at *attB*, the transcription direction of *dnaAp1* and *dnaAp2* promoters is the same as movement of the replication fork. This orientation is the same as at the *dnaA* wild type location. The genetic location of the fusions was confirmed in the *Cm<sup>r</sup>* transformants by demonstrating cotransduction of *Cm<sup>r</sup>* and *galK*.

15

The fusions created in strain WM2269 were moved by P1 transduction to MG1655, creating strain DYK9W with pDYK9 and strain DYK11W with pDYK11. Deletion of the *rpmH* promoters had very little effect (1.4-fold) on chitobiase activity (Table 2, compare lines 1 and 3)

20

**Table 2.**  
**Chitobiase and  $\beta$ -Gal Activities of *dnaA-chb* and *dnaA-lacZ* Fusions in WT and *fis* Mutant Backgrounds**

Strains	$\beta$ -Gal <sup>c</sup>	
	Chitobiase <sup>a</sup> (U <sup>b</sup> )	(Miller U <sup>d</sup> )
DYK11W <i>fis</i> +	30.0 $\pm$ 2.1	
DYK11F <i>fis</i> ::985	80.0 $\pm$ 1.2	
DYK9W <i>fis</i> +	44.0 $\pm$ 2.1	
DYK9F <i>fis</i> ::985	96.0 $\pm$ 3.5	
RB220 <i>fis</i> +		59.8 $\pm$ 7.9
TP220 <i>fis</i> ::767		115.2 $\pm$ 7.2
<sup>a</sup> Triplicate samples were assayed during exponential growth. Mean chitobiase activities are given with standard deviation. <sup>b</sup> One unit of chitobiase activity is 1 pmol of <i>p</i> -nitrophenol per min at 28°C. Units given for 1 mL of culture at OD <sub>450</sub> = 1. <sup>c</sup> Data from Reference 8. <sup>d</sup> Unit defined in Reference 13.		

Fis protein binds to a site in the *dnaA*<sub>P2</sub> promoter that covers the -35 sequence (Figure 5), and appears to be a repressor of DnaA expression. A fusion protein with  $\beta$ -galactosidase activity that is expressed from the *rpmH-dnaA* regulatory region has increased  $\beta$ -galactosidase activity (1.9-fold) in a *fis*<sup>-</sup> mutant when compared to Fis wild type cells [as described in Froelich, J. M., et al., "Fis binding in the *dnaA* operon promoter region," J. Bacteriol. 178:6006-6012 (1996), data shown in Table 3]. Similarly, the absence of Fis leads to a greater than 2-fold increase in chitobiase activity of the DnaA-chitobiase fusion protein for the DYK9F and DYK11F strains, comparable in extent to that observed with the *dnaA-lacZ* fusion strain, TP220 (Table 3).

In the reporter gene constructs discussed in the Example, all upstream transcriptional activity was prevented from entering the *chb* reporter gene. The plasmid vectors, pJMF3 and pJMF4, described in the Examples above, contained the *rmbt1t2* terminator upstream of the promoter fusion, which prevented readthrough from chromosomal promoters near the insertion site. Only chitobiase activity originating from the promoters of interest was expressed.

The *attP* site in these vectors allowed integration at the chromosomal *attB* in a specific orientation, depending on the vector used. With these vectors, any chitobiase fusion involving an essential gene can be moved to the chromosome, thus permitting single copy analysis with a chromosomal orientation similar to the wild-type gene.

#### Example 7

##### Identification of Promoters in Test Sequences

A nucleic acid prospectively containing a promoter is inserted upstream of a nucleic acid encoding a cytoplasmic form of chitobiase as described above. For example, the nucleic acid prospectively containing a promoter may be inserted into a restriction site in a sequence containing a plurality of restriction sites, such as a polylinker, which is located upstream of the nucleic acid encoding chitobiase. The test sequence may comprise any nucleic acid to be tested for promoter activity. In one embodiment, the test sequence may comprise a genomic DNA sequence. For example, the genomic DNA sequence may be a randomly generated DNA fragment, such as a fragment generated using shotgun cloning techniques, a restriction fragment, or any other sequence.

The vectors containing the test sequence upstream of the nucleic acid encoding chitobiase are introduced into an appropriate host cell. The level of chitobiase activity is assayed and compared to the level obtained from a control vector which lacks an insert in the cloning site. The presence of an elevated expression level in cell containing the vector containing the insert with respect to the level in cells containing the control vector without the insert indicates the presence of a promoter in the insert.

In some embodiments, the activity of the promoter in the test sequence may be assayed after exposure of the host cells to conditions which may influence the level of transcription from the promoter. For example, the environment of the host cells may be altered to determine whether the transcription level is influenced by environmental factors, including factors such as temperature, pH, nutrients, or availability of oxygen. In such analyses, chitobiase levels are assayed under a variety of environmental conditions to determine the effects of the environmental conditions on transcription levels from the promoter. In addition, the activity of the promoters may be examined in the presence or absence of compounds to be



tested for regulatory activity. For example, the activity of the promoters may be tested by determining the levels of chitobiase produced in the presence or absence of compounds to be tested for activity as drugs.

Promoter sequences within the test sequences may be further defined by constructing nested deletions in the test sequences using conventional techniques such as Exonuclease III digestion. The resulting deletion fragments can be inserted into the promoter reporter vector to determine whether the deletion has reduced or obliterated promoter activity as determined by measuring chitobiase activity in cells containing the deletion vectors. In this way, the boundaries of the promoters may be defined. If desired, potential individual regulatory sites within the promoter may be identified using techniques such as site directed mutagenesis or linker scanning to obliterate potential transcription factor binding sites within the promoter individually or in combination. The effects of these mutations on transcription levels may be determined by inserting the mutations into the cloning sites in the promoter reporter vectors and measuring the levels of chitobiase produced from the mutated promoters.

The activity of known promoters may also be monitored by operably linking them to a nucleic acid encoding a cytoplasmic form of chitobiase. The activity of the promoters may be analyzed under various environmental conditions as described above. In addition, the activity of the promoters may be analyzed in the presence or absence of compounds to be tested for the ability to affect transcription from the promoters. For example, the compounds may be tested for activity as drugs.

In some embodiments, the chitobiase reporter constructs may be used in systems for identifying compounds that modulate cell surface protein-mediated activity or compounds which modulate the activities of intracellular signaling systems. Techniques for using reporter genes to identify compounds which modulate cell surface protein-mediated activity have been described in U.S. Patent Number 5,401,629 and U.S. Patent Number 5,436,128. Briefly, in such methods, a construct comprising a promoter operably linked to a nucleic acid encoding a reporter enzyme is introduced into cells which express the cell surface protein and cells which do not express the cell surface protein. Each of the cells are contacted with test compounds and the effects of these compounds on transcription levels is measured by determining the level of activity of the reporter enzyme. The level of expression of the reporter gene in cells expressing the cell surface protein is compared to the level in cells which do not express the cell surface protein to identify compounds that modulate cell surface protein activity.

Similarly, the chitobiase reporter constructs may be used to identify compounds which influence the activity of intracellular signaling pathways, such as cAMP-based or phosphorylation-based pathways. In such methods, a promoter which is activated via such pathways is operably linked to a nucleic acid encoding a cytoplasmic form of chitobiase. The cells are contacted with test compounds. Those compounds which activate the pathway to which the promoter responds will produce an enhanced level of chitobiase activity in the cells as compared to the level of chitobiase activity in control cells which have not been contacted with the test compound.

## Example 8

Detecting Successful Transformation or Transfection Using Chitobiase

5 A vector comprising a sequence encoding a cytoplasmic form of chitobiase operably linked to a sequence capable of directing transcription of the chitobiase gene is introduced into a host cell. The host cells are contacted with a chitobiase substrate and those host cells which contain chitobiase activity are identified as cells which were successfully transformed or transfected. In some embodiments, a portion or replica of a colony may be lysed or permeabilized prior and the lysate or permeabilized cells may be contacted with the chitobiase substrate.

## CONCLUSION

10 New gene reporter systems that use chitobiase have been described. Chitobiase has advantages over other reporter gene systems in that chitobiase is not found in many cell lines traditionally used in reporter gene systems.

Finally, the forgoing examples are not intended to limit the scope of the present invention, which is set forth in the following claims. In particular, various equivalents and substitutions will be recognized by those of ordinary skill in the art in view of the foregoing disclosure, and these are contemplated to be within the scope of the present invention.

15

WHAT IS CLAIMED IS:

1. A method for characterizing a promoter comprising:
  - providing a construct comprising said promoter operably linked to a nucleic acid encoding a cytoplasmic form of chitobiase;
  - 5           introducing the construct into host cells; and
  - identifying sequences in said promoter which regulate transcription levels.
2. The method of Claim 1, wherein said cytoplasmic form of chitobiase lacks a signal sequence.
3. The method of Claim 2, wherein said nucleic acid encoding a cytoplasmic form of chitobiase encodes a fusion protein, said fusion protein comprising a cytoplasmic form of chitobiase fused to a heterologous polypeptide.
- 10       4. The method of Claim 1, wherein said nucleic acid encoding a cytoplasmic form encodes a cytoplasmic form of chitobiase obtained from an organism selected from the group consisting of *Alteromonas sp. 0-7*, *Arabidopsis thaliana*, *Bacillus subtilis*, *Bombyx mori*, *Bos taurus*, *Caenorhabditis elegans*, *Candida albicans*, *Dictyostelium discoideum*, *Entamoeba histolytica*, *Felis catus*, *Homo sapiens*, *Korat cats*, *Lactobacillus casei*, *Leishmania donovani*, *Mus musculus*, *Pisum sativum*, *Porphyromonas gingivalis*, *Pseudoalteromonas sp. S9*, *Rattus norvegicus*, *Serratia marcescens*, *Streptomyces plicatus*, *Streptomyces thermoviolaceus*, *Sus scrofa*, *Trichoderma harzianum*, *Vibrio furnissii*, *Vibrio harveyi*, *Vibrio parahaemolyticus*, and *Vibrio vulnificus*.
- 15       5. The method of Claim 1, wherein said method of identifying sequences which are involved in directing transcription comprises mutagenizing said promoter.
6. The method of Claim 1, wherein said method of identifying sequences which are involved in
- 20       transcription comprises constructing deletions in said promoter.
7. A method for identifying a regulatory element capable of directing or regulating transcription within a test nucleic acid sequence comprising:
  - providing a construct comprising said test nucleic acid sequence operably linked to a nucleic acid encoding a cytoplasmic form of chitobiase;
  - 25           introducing said construct into host cells; and
  - determining the level of chitobiase activity.
8. The method of Claim 7, wherein said cytoplasmic form of chitobiase lacks a signal sequence.
9. The method of Claim 8, wherein said nucleic acid encoding a cytoplasmic form of chitobiase encodes a fusion protein, said fusion protein comprising a cytoplasmic form of chitobiase fused to a heterologous polypeptide.
- 30       10. The method of Claim 7, wherein said nucleic acid encoding a cytoplasmic form encodes a cytoplasmic form of chitobiase obtained from an organism selected from the group consisting of *Alteromonas sp. 0-7*, *Arabidopsis thaliana*, *Bacillus subtilis*, *Bombyx mori*, *Bos taurus*, *Caenorhabditis elegans*, *Candida albicans*, *Dictyostelium discoideum*, *Entamoeba histolytica*, *Felis catus*, *Homo sapiens*, *Korat cats*, *Lactobacillus casei*, *Leishmania donovani*, *Mus musculus*, *Pisum sativum*, *Porphyromonas gingivalis*, *Pseudoalteromonas sp. S9*, *Rattus norvegicus*, *Serratia*

*marcescens*, *Streptomyces plicatus*, *Streptomyces thermoviolaceus*, *Sus scrofa*, *Trichoderma harzianum*, *Vibrio furnissii*, *Vibrio harveyi*, *Vibrio parahaemolyticus*, and *Vibrio vulnificus*.

11. The method of Claim 7, wherein said reporter gene construct is introduced transiently.
12. The method of Claim 7, wherein said reporter gene construct is introduced stably.
- 5 13. The method of Claim 7, wherein said host cells are selected from the group consisting of prokaryotic cells and eukaryotic cells.
14. The method of Claim 7, further comprising permeabilizing or lysing said host cells.
15. The method of Claim 14, wherein said permeabilizing or lysing step comprises treating said host cells with toluene.
- 10 16. The method of Claim 7, wherein said step of determining the level of chitinase activity is selected from the group consisting of measuring the amount of a chemiluminescent product produced from a substrate, measuring the amount of a fluorescent product produced from a substrate, measuring the amount of light absorbed by a product produced from a substrate and measuring a decrease in the amount of a detectable substrate.
17. The method of Claim 7, wherein said step of determining the level of chitinase activity comprises
  - 15 determining the level of *p*-nitrophenol released from a substrate.
  18. The method of Claim 7, wherein said test nucleic acid sequence comprises a portion of genomic DNA.
  19. The method of Claim 7, wherein said step of determining the level of chitinase activity comprises determining the level of chitinase activity after exposing said host cells to a desired set of environmental conditions.
  20. The method of Claim 7, wherein said step of determining the level of chitinase activity comprises
    - 20 determining the level of chitinase activity after contacting said host cells with a compound to be tested for its influence on the level of transcription from said regulatory element.
    21. A method of detecting successful transformation, comprising the steps of:
      - introducing a nucleic acid encoding a cytoplasmic form of chitinase into host cells; and
      - detecting chitinase expression in said host cells.
    - 25 22. A fusion protein-reporter gene construct comprising a promoter operably linked to a nucleic acid encoding a cytoplasmic form of chitinase fused in frame with a nucleic acid encoding a heterologous polypeptide, wherein said heterologous polypeptide is not -galactosidase or a portion thereof and wherein said heterologous polypeptide does not contain a signal peptide.
    23. The nucleic acid of Claim 22, wherein said nucleic acid encodes a cytoplasmic form of chitinase
      - 30 obtained from an organism selected from the group consisting of *Alteromonas sp. O-7*, *Arabidopsis thaliana*, *Bacillus subtilis*, *Bombyx mori*, *Bos taurus*, *Caenorhabditis elegans*, *Candida albicans*, *Dictyostelium discoideum*, *Entamoeba histolytica*, *Felis catus*, *Homo sapiens*, *Korat cats*, *Lactobacillus casei*, *Leishmania donovani*, *Mus musculus*, *Pisum sativum*, *Porphyromonas gingivalis*, *Pseudoalteromonas sp. S9*, *Rattus norvegicus*, *Serratia marcescens*, *Streptomyces plicatus*, *Streptomyces thermoviolaceus*, *Sus scrofa*, *Trichoderma harzianum*, *Vibrio furnissii*, *Vibrio harveyi*, *Vibrio parahaemolyticus*, and *Vibrio vulnificus*.

24. The nucleic acid of Claim 22, further comprising a  $\lambda$  site-specific recombination sequence.
25. A reporter gene construct comprising plasmid pJMF3.
26. A reporter gene construct comprising plasmid pJMF4.
27. A reporter gene construct comprising plasmid pDYK9.
- 5 28. A reporter gene construct comprising plasmid pDYK11.
29. A host cell comprising the construct of Claim 22.
30. The host cell of Claim 29 wherein said nucleic acid is integrated into a chromosome of said cell.
31. The host cell of Claim 29, wherein said nucleic acid is transiently expressed in said host cell.
32. A nucleic acid encoding a cytoplasmic form of chitobiase in which the signal sequence of native  
10 chitobiase has been inactivated or deleted.
33. The nucleic acid of Claim 32, wherein the signal sequence has been mutated to inactivate it.
34. An isolated or purified polypeptide comprising a cytoplasmic form of chitobiase fused in frame with a  
heterologous polypeptide, wherein said heterologous polypeptide is not -galactosidase or a portion thereof and  
wherein said heterologous polypeptide does not contain a signal peptide.
- 15 35. An isolated or purified polypeptide comprising a cytoplasmic form of chitobiase in which the signal  
peptide of native chitobiase has been inactivated or deleted.
36. The polypeptide of Claim 35, wherein the signal sequence has been mutated to inactivate it.
37. A method for monitoring the activity of a promoter comprising:  
providing a construct comprising said promoter operably linked to a nucleic acid encoding a  
20 cytoplasmic form of chitobiase;  
introducing said construct into host cells; and  
determining the level of chitobiase activity.
38. The method of Claim 37, wherein said cytoplasmic form of chitobiase lacks a signal sequence.
39. The method of Claim 38, wherein said nucleic acid encoding a cytoplasmic form of chitobiase encodes a  
25 fusion protein, said fusion protein comprising a cytoplasmic form of chitobiase fused to a heterologous polypeptide.
40. The method of Claim 37, wherein said nucleic acid encoding a cytoplasmic form encodes a cytoplasmic  
form of chitobiase obtained from an organism selected from the group consisting of *Alteromonas sp. O-7*, *Arabidopsis*  
*thaliana*, *Bacillus subtilis*, *Bombyx mori*, *Bos taurus*, *Caenorhabditis elegans*, *Candida albicans*, *Dictyostelium*  
*discoideum*, *Entamoeba histolytica*, *Felis catus*, *Homo sapiens*, *Korat cats*, *Lactobacillus casei*, *Leishmania donovani*,  
30 *Mus musculus*, *Pisum sativum*, *Porphyromonas gingivalis*, *Pseudoalteromonas sp. S9*, *Rattus norvegicus*, *Serratia*  
*marcescens*, *Streptomyces plicatus*, *Streptomyces thermoviolaceus*, *Sus scrofa*, *Trichoderma harzianum*, *Vibrio*  
*furnissii*, *Vibrio harveyi*, *Vibrio parahaemolyticus*, and *Vibrio vulnificus*.
41. The method of Claim 37, wherein said reporter gene construct is introduced transiently.
42. The method of Claim 37, wherein said reporter gene construct is introduced stably.

43. The method of Claim 37, wherein said host cells are selected from the group consisting of prokaryotic cells and eukaryotic cells.

44. The method of Claim 37, further comprising permeabilizing or lysing said host cells.

5 45. The method of Claim 44, wherein said permeabilizing or lysing step comprises treating said host cells with toluene.

46. The method of Claim 37, wherein the step of determining the level of chitinase activity is selected from the group consisting of measuring the amount of a chemiluminescent product produced from a substrate, measuring the amount of a fluorescent product produced from a substrate, measuring the amount of light absorbed by a product produced from a substrate and measuring a decrease in the amount of a detectable substrate.

10 47. The method of Claim 37, wherein said step of determining the level of chitinase activity comprises determining the level of *p*-nitrophenol released from a substrate.

48. The method of Claim 37, wherein said step of determining the level of chitinase activity comprises determining the level of chitinase activity after exposing said host cells to a desired set of environmental conditions.

15 49. The method of Claim 37, wherein said step of determining the level of chitinase activity comprises determining the level of chitinase activity after contacting said host cells with a compound to be tested for its influence on the level of transcription from said regulatory element.

50. The method of Claim 49, wherein said compound comprises a compound to be tested for activity as a drug.




Figure 1

SEQ ID NO: 15:




Figure 2








Figure 3

**Figure 4**



SEQ ID NO: 17

Figure 5

## SEQUENCE LISTING

<110> ELITRA PHARMACEUTICALS, INC.  
Zyskind, Judith W.

<120> CHITOBIASE AS A REPORTER ENZYME

<130> ELITRA.006VPC

<150> 60/159,221

<151> 1999-10-13

<160> 19

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide

<400> 1

gcacatgcat gctgggcatt aaattttcc

29

<210> 2

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide

<400> 2

cggggtacca actcatcctg caatcg

26

<210> 3

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide

<400> 3

ctagtctaga tgccgaactc agaagtga

28

<210> 4

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide

<400> 4

gcacatgcat gcgggggatg gctttagat

30

<210> 5

<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide

<400> 5  
catgatcatg cgacagggtt gatga

25

<210> 6  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide

<400> 6  
gggggcgcct acctttcacg ag

22

<210> 7  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide

<400> 7  
catgcatgca tgaaacgatg gacacc

26

<210> 8  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide

<400> 8  
cggggtacca actcatcctg caatcg

26

<210> 9  
<211> 12  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide

<400> 9  
cattaatgca tg

12

<210> 10  
<211> 12  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Oligonucleotide

<400> 10  
cattaatgca tg

12

<210> 11  
<211> 5826  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> pDYK9

<400> 11  
gaattccgga tgagcattca tcaggcgggc aagaatgtga ataaaggccg gataaaactt 60  
gtgcttattt ttctttacgg tctttaaaaa ggccgtaata tccagctgaa cggctctggtt 120  
ataggtacat tgagcaactg actgaaatgc ctcaaaatgt tctttacgat gccattggga 180  
tatatcaacg gtggtatata cagtgatatt tttctccatt ttagcttccg tagctcctga 240  
aaatctcgat aactcaaaaa atacgcccgg tagtgatctt atttcattat ggtgaaagtt 300  
ggaacctctt acgtgccgat caacgtctca ttttcgcaa aagttggccc agggcttccc 360  
ggtatcaaca gggacaccag gatttattta ttctgcgaag tgatcttccg tcacagggtat 420  
ttattcggcg caaagtgcgt cgggtgatgc tgccaactta ctgatttagt gtatgatggt 480  
gtttttgagg tgctccagtg gcttctgttt ctatcagctg tccctcctgt tcagctactg 540  
acggggtggt cgctaaccgg aaaagcaccg ccggacatca gcgctagcgg agtgtgcggc 600  
cgcatggct tactatgttg gcaactgatga ggggtgcagt gaagtgtctt atgtggcagg 660  
agaaaaaagg ctgcaccggt gcgtcagcag aatatgtgat acaggatata ttccgcttcc 720  
tcgctcactg actcgctacg ctccgtcggt cgactgcggc gagcggaaat ggcttacgaa 780  
cggggcggag atttcctgga agatgccagg aagatactta acagggaagt gagaggccg 840  
cggcaaagcc gtttttccat aggtccggcc ccctgacaa gcatcacgaa atctgacgct 900  
caaactcagt gtggcgaaac ccgacaggac tataaagata ccaggcggtt ccctggcg 960  
ctccctcggt cgctctcctg ttctgcctt cgggtttacc ggtgtcattc cgctgttatg 1020  
gccgcgtttg tctcattcca cgcctgacac tcagttccgg gtaggcagtt cgctccaagc 1080  
tggaactgtat gcacgaaccc ccggttcagt ccgaccgctg cgccttatcc ggtaactatc 1140  
gtcttgagtc caaccoggaa agacatgcaa aagcaccact ggcagcagcc actggttaatt 1200  
gatttagagg agttagtctt gaagtcatgc gccggttaag gctaaactga aaggacaagt 1260  
tttgggtgact gcgtcctctc aagccagtta cctcggttca aagagttggt agctcagaga 1320  
accttcgaaa aaccgcccgt caaggcgggt ttttcgtttt cagagcaaga gattacgcgc 1380  
agaccaaaac gatctcaaga agatcatctt atgcggccgc atcagataaa atatttctag 1440  
atgccgaact cagaagtga acgccgtagc gccgatggta gtgtggggtc tccccatgcg 1500  
agagtaggga actgccaggc atcaaataaa acgaaaggct cagtcgaaag actgggcctt 1560  
tcgttttatc tgttgtttgt cgtgaaacgc tctctgagt aggacaaatc cgcggggagc 1620  
ggatttgaac gttgcgaagc aacggcccgg aggggtggcg gcaggacgcc cgccataaac 1680  
tgccaggcat caaattaagc agaaggccat cctgacggat ggcctttttg cgtttctaca 1740  
aactcttctc gctgcatat ctacaagcca tccccgcgt gctggtcatt aaattttcca 1800  
atatgcggcg taaatcgtgc ccgcctcgcg gcaggtcgt ttacacttag cgagttctgg 1860  
aaagtcctgt ggataaatcg ggaatctgt tgagaaacag aagatctctt gcgcagttta 1920  
ggctatgata cgcggtcccg atcggtttgc aggatcttga tcgggcatat aaccgcagac 1980  
agcgggtcgt gcgtcaccct caagcagggt ctttctgacg tacgtcaaca atcatgaatg 2040  
tttcagcctt agtcattatc gacttttgtt cgagtgaggt ccgccgtgtc actttcgtt 2100  
tggcagcagt gtcttgcccg attgcaggat gatttggtac cgagctctga gcaacaagtt 2160  
gtaaactcac tggctgataa ccttgatata caatatgaag tgtaactaa ccatggtgct 2220  
aacgaaggct ttgcgtgcca agatatgggc gcagaatggg cttctgttaa caaagttaac 2280  
atgacgctt ttaaccaagg tgaagctgtt gactcaaaag attgggctat ttacttccac 2340  
agcattcgtc tgattctgga tgttgacaac gagcagttca aaatctctcg tgtaacgggt 2400  
gacctacata agctagaacc aacagataag tttgacggct tcgctgcggg tgaagaggtt 2460  
gttcttccat tgggttggtga atactggcaa ctatttgaaa ctgacttcat gccgggtgca 2520  
ttcgtttctg ctccaaacgc agaacctaa atgattgctt ctctaaatac tgaagatgtt 2580  
gcgtcttttg tgacgggtct tgaaggtaac aaactaaac gtacaccaga tgacaacaat 2640  
gtattttcaa acgtgtgtc tcgttttgag aaaaacgaag acctagcaac acaagacgta 2700  
tcaaccacgt tactaccaac accaatgcac gttgaagcgg gtaaaaggcaa agtagatata 2760  
gcggatggta ttgcgtgcc taaagacgca ttcgatgcga ctcagttcgc agcgattcaa 2820  
gatcgtgcag aagtggtagg tgtggacgtt cgtggtgatc ttctgtaag catcactgtt 2880

gttcctgcag	acttcaccgg	tgaattagca	aaatctggtg	cttacgaaat	gagcatcaaa	2940
ggcgacggta	ttgtgattaa	agcggttcgac	caagcaggcg	ctttctacgc	agtacaatct	3000
atctttggcc	tggtagatag	ccaaaatgct	gattctctac	cacaactgtc	tattaaagat	3060
gcgcctcggt	ttgattaccg	tggtgtgatg	gtggatgtgg	ctcgtaactt	ccactctaag	3120
gacgcaatcc	ttgcaacgct	agaccaaagt	gcagcgtaaca	agatgaacaa	acttcacctt	3180
cacctaaccg	atgatgaagg	ctggcggtta	gaaatcccgg	gtctgcctga	gctgacagaa	3240
gtgggtgcta	accgttggtt	cgatacacia	gagaaaagct	gtttactgcc	tcagcttggc	3300
tctggtcaaa	cgacagacaa	ctttggctct	ggctacttca	gcaaagcaga	ctacgtggaa	3360
atcttgaaat	acgcgaaagc	acgtaacatt	gaagtgtatt	cagaaatcga	tatgccagct	3420
cacgctcggt	cagcagtagt	atcaatggaa	gctcggttacg	accgcctaata	ggaagaaggt	3480
aaagaagctg	aagcgaacga	ataccgtctg	atggatcctc	aagatacatc	aaacgtaacg	3540
acggttcagt	tctacaataa	gcaaagcttc	atcaacccat	gtatggaatc	ttcaactcgc	3600
tttggtgata	aggtgatttc	agaagtggca	gcaatgcacc	aagaagctgg	cgctccacta	3660
acaacttggc	acttcgggtg	tgacgaagcg	aagaacatca	agctagggtg	tggtttccaa	3720
gacgttaacg	cagaagataa	agtaagctgg	aaaggcacga	ttgacctgtc	ttaacaagac	3780
aagccgtttg	cacagtctcc	acaatgtcag	acgctaatac	cagatggcac	agtcagtgc	3840
tttgctcacc	taccaagcca	cttcgcggaa	gaagtgtcga	agattgttgc	tgagaaaggc	3900
attccaaact	tccaagcttg	gcaagatggt	ttgaaatata	gtgacggcga	aaaagcggtc	3960
gctacagaaa	atactcgcgt	aaacttctgg	gacgttctgt	actggggcgg	tacttcctca	4020
gtgtacgagt	ggtctaagaa	aggttacgac	gtgattgttt	ctaaccacaga	ttacgtgtac	4080
atggatatgc	catacgaagt	tgacccgaaa	gagcgtgggt	actactgggc	aacacgtgca	4140
acggatactc	gtaagatgtt	tggttttgca	ccaggaacaa	tgacctcaaaa	cgcagaaact	4200
tctgtagatc	gcgatggcaa	tggttttact	ggtaaagggtg	aaatcgaagc	gaaacctttc	4260
tacggctctat	ctgcacaact	ttgggtctgag	acagtacgta	acgacgagca	atagcagtag	4320
atggatttcc	ctcgcgtcct	cgctgctgct	cagcgtgcat	ggcaccgtgc	tgactgggaa	4380
aacgactaca	aagttgggtg	tgagtactcg	caaaactcta	atctagttga	taaagcatcg	4440
ctaaaccaag	actacaaccg	ctttgcgaac	gtacttggtc	aacgtgaact	ggctaagcta	4500
gaaaaatcag	gtattgacta	cgcctaccac	gtaccagggtg	caaaagtaga	agatggtaag	4560
ctagcaatga	acgttcagtt	ccctggcgta	acgcttcaat	actctctgga	tggtgagaac	4620
tggttgactt	atgcagacaa	cgctcgtcca	aatgtaactg	gtgaagtctt	catccgctcg	4680
gtatctgcga	caggtgagaa	ggtaagccgt	atcactagcg	tgaaataata	gcgctcagta	4740
ttcactaaaa	tcatagtctc	ttactcaaa	ccctcaactt	atgttggggg	ctttgtttat	4800
ttttcttcgg	aaaataagcg	tgatcatcg	acagggttga	tgacaaaaaa	ttagcgcaag	4860
aagacaaaaa	tcaccttgcg	ctaagtctct	gttacagggtc	actaatacca	tctaagtagt	4920
tgattcatag	tgactgcata	tggtgtgttt	tacagtatta	tgtagtctgt	tttttatgca	4980
aaatctaatt	taatatattg	atattttatat	cattttacgt	ttctcgttca	gcttttttat	5040
actaagttgg	cattataaaa	aagcattgct	tatcaatttg	ttgcaacgaa	caggtcacta	5100
tcagtcaaaa	taaaatcatt	atgttgatttc	aattttgtcc	cactccctgc	ctctgtcatc	5160
acgatactgt	gatgccatgg	tgtccgactt	atgcccagaa	agatgttgag	caaacttatc	5220
gcttatctgc	ttctcataga	gtcttgacga	caaactgcgc	aactcgtgaa	aggtaggcgc	5280
ccccggtcga	atgttgcttc	gaatttctgc	cattcatccg	cttattatca	cttattcagg	5340
cgtagcacca	ggcgtttaag	ggcaccaata	actgccttaa	aaaaattacg	ccccgcctg	5400
ccactcatcg	cagtactggt	gtaattcatt	aagcattctg	ccgacatgga	agccatcaca	5460
gacggcatga	tgaacctgaa	tcgccagcgg	catcagcacc	ttgtcgcctt	gcgtataata	5520
tttgcccatg	gtgaaaacgg	gggcgaagaa	gttgtccata	ttggccacgt	ttaaatcaaa	5580
actggtgaaa	ctcaccagg	gattggctga	gacgaaaaac	atattctcaa	ttaacccttt	5640
agggaaatag	gccagggttt	caccgtaaca	cgccacatct	tgcgaaatata	tgtgtagaaa	5700
ctgccggaaa	tcgtcgtggt	attcactcca	gagcgatgaa	aacgtttcag	tttgctcatg	5760
gaaaacgggtg	taacaagggt	gaacactatc	ccatatacacc	agctcaccgt	ctttcattgc	5820
catacg						5826

&lt;210&gt; 12

&lt;211&gt; 6071

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; pDYK11

&lt;400&gt; 12

gaattccgga tgagcattca tcaggcgggc aagaatgtga ataaaggccg gataaaactt

60

gtgcttattt	ttcttttacgg	tcttttaaaaa	ggccgtaata	tccagctgaa	cggtctgggt	120
ataggtacat	tgagcaactg	actgaaatgc	ctcaaaatgt	tctttacgat	gccattggga	180
tatatcaacg	gtggtatata	cagtgatatt	tttctccatt	ttagcttcct	tagctcctga	240
aaatctcgat	aactcaaaaa	atacgcccg	tagtgatctt	atttcattat	ggtgaaagtt	300
ggaacctctt	acgtgccgat	caacgtctca	ttttcgccaa	aagttggccc	agggtctccc	360
ggtatcaaca	gggacaccag	gatttattta	ttctgcgaag	tgatcttccg	tcacagggtat	420
ttattcggcg	caaagtgcgt	cggtgatgc	tgccaactta	ctgatttagt	gtatgatggt	480
gtttttgagg	tgctccagt	gcttctgttt	ctatcagctg	tccctcctgt	tcagctactg	540
acgggggtgt	gcgtaacggc	aaaagcaccg	ccggacatca	gcgctagcgg	agtgtgcggc	600
cgcactgggt	tactatgttg	gcactgatga	gggtgtcagt	gaagtgcctc	atgtggcagg	660
agaaaaaagg	ctgcaccggg	gcgtcagcag	aatatgtgat	acaggatata	ttccgcttcc	720
tcgctcactg	actcgctacg	ctcggctcgt	cgactgcggc	gagcggaaat	ggcttacgaa	780
cggggcgagg	atttcctgga	agatgccagg	aagatactta	acagggaagt	gagagggccg	840
cggcaaagcc	gtttttccat	aggtccggcc	cccctgacaa	gcatacagaa	atctgacgct	900
caaatcagtg	gtggcgaaac	ccgacaggac	tataaagata	ccaggcggtt	cccctggcgg	960
ctccctcgtg	cgtctcctg	ttcctgcctt	tcgggtttacc	ggtgtcattc	cgctgttatg	1020
gccgcgtttg	tctcattcca	cgctgcacac	tcagttccgg	gtaggcagtt	cgctccaagc	1080
tggactgtat	gcacgaaccc	ccggttcagt	ccgaccgctg	cgcttattcc	ggtaactatc	1140
gtcttgatgc	caaccgcgaa	agacatgcaa	aagcaccact	ggcagcagcc	actggttaatt	1200
gatttagagg	agtttagtctt	gaagtcatgc	gccgggttaa	gctaaactga	aaggacaagt	1260
tttggtagct	gcgtcctccc	aagccagtta	cctcgggtta	aagagttggt	agctcagaga	1320
accctcgaaa	aaccgcccgt	caaggcggtt	tttctgtttt	cagagcaaga	gattacgcgc	1380
agaccaaaac	gatctcaaga	agatcatctt	atgcggccgc	atcagataaa	atatttctag	1440
atgccgaact	cagaagtga	acgccgtagc	gccgatggta	gtgtggggtc	tcccatgcg	1500
agagtaggga	actgccaggc	atcaaataaa	acgaaaggct	cagtcgaaag	actgggcctt	1560
tcgttttata	tggtgtttgt	cgggtgaacgc	tctcctgagt	aggacaaatc	cgccgggagc	1620
ggatttgaac	gttgcgaagc	aacggcccgg	aggggtggcg	gcaggacgcc	cgccataaac	1680
tgccaggcat	caaatgaagc	agaaggccat	cctgacggat	ggcctttttg	cgtttctaca	1740
aactcttctt	gtcgtcatat	ctacaagcca	tccccgcgat	gcatgaaacg	atggacaccg	1800
aagccatggg	tgattaaaga	ggccggattg	taataattgt	acactccgga	gtcaattctc	1860
tttcttattt	taccgcgctt	ttccgcacct	tttcgcaggg	aaaatgtacg	acctcacacc	1920
agtggaaacc	agcatggcgc	gccgggtgga	ggattatacg	ggctgatggg	taaagcgcaa	1980
ggatcgtcct	ggatctttat	tagatcgatt	aagccaattt	ttgtctatgg	tcattaaatt	2040
ttccaatatg	cggcgtaaat	cgtgcccgcc	tcgcggcagg	atcgtttaca	cttagcgagt	2100
tctggaagt	cctgtggata	aatcgggaaa	atctgtgaga	aacagaagat	ctcttgcgca	2160
gtttaggcta	tgatccgcgg	tcccgcgt	tttgcaggat	cttgatcggg	catataaccg	2220
cagacagcgg	ttcgtgcgtc	accctcaagc	agggctcttt	cgacgtacgt	caacaatcat	2280
gaatgtttca	gccttagtca	ttatcgactt	ttgttcgagt	ggagtccgcc	gtgtcacttt	2340
cgctttggca	gcagtgtctt	gcccgaattg	aggatgagtt	ggtaccgagc	tctgagcaac	2400
aagttgtaaa	ctcactggct	gataaccttg	atatccaata	tgaagtgtta	actaaccatg	2460
gtgctaacga	aggtcttgcg	tgccaagata	tgggcgcgaga	atgggcttct	tgtaacaaag	2520
taaacatgac	gcttgttaac	caagggtgaag	ctgttgactc	aaaagattgg	gctatttact	2580
tccacagcat	tcgtctgatt	ctggatgttg	acaacgagca	gttcaaaatc	tctcgtgtaa	2640
cgggtgacct	acataagcta	gaaccaacag	ataagtttga	cggttctcgt	gccggtgaag	2700
aggttgttct	tccattgggt	ggtgaatact	ggcaactatt	tgaaactgac	ttcatgccgg	2760
gtgcattcgt	ttctgctcca	aacgcagaac	ctaagatgat	tgcttctcta	aatactgaag	2820
atgttgcgtc	ttttgtgacg	ggtcttgaag	gtaacaacct	aaaacgtaca	ccagatgaca	2880
acaatgtatt	tgcaaacgct	gtgtctcgtt	ttgagaaaaa	cgaagacctc	gcaacacaag	2940
acgtatcaac	cacgttacta	ccaacaccaa	tgcacgttga	agcgggtaaa	ggcaaagtag	3000
atatcgcgga	tggtattgcg	ctgcctaaag	acgcattcga	tgcgactcag	ttcgcagcga	3060
ttcaagatcg	tgcaagaatg	gtagggtgtg	acgttctgtg	tgatcttccct	gtaagcatca	3120
ctgttgttcc	tgcaagactt	accggtgaat	tagcaaaatc	tggtgcttac	gaaatgagca	3180
tcaaaggcga	cggattatgt	attaaagcgt	tcgaccaagc	aggcgctttc	tacgcagtag	3240
aatctatctt	tggcctggta	gatagccaaa	atgctgattc	tctaccacaa	ctgtctatta	3300
aagatgcgcc	tcgttttgat	taccgtgggt	tgatgggtga	tgtggctcgt	aacttccact	3360
ctaaggacgc	aatccttgca	acgctagacc	aaatggcagc	gtacaagatg	aacaaacttc	3420
accttcacct	aaccgatgat	gaaggctggc	gtttagaaat	cccggtctcg	cctgagctga	3480
cagaagtggg	tgctaaccgt	tgtttcgata	cacaagagaa	aagctgttta	ctgcctcagc	3540
ttggctctgg	tccaacgaca	gacaactttg	gctctggcta	cttcagcaaa	gcagactacg	3600
tggaatctt	gaaatacgcg	aaagcacgta	acattgaagt	gattccagaa	atcgatatgc	3660
cagctcacgc	tcgtgcagca	gtagtatcaa	tggaagctcg	ttacgaccgc	ctaagtgaag	3720



```

aaggtaaaga agctgaagcg aacgaatacc gtctgatgga tcctcaagat acatcaaacg 3780
taacgcagggt tcagttctac aataagcaaa gtttcatcaa cccatgtatg gaatcttcaa 3840
ctcgctttgt tgataagggtg atttcagaag tggcagcaat gcaccaagaa gctggcgctc 3900
cactaacaac ttggcacttc ggtggtgacg aagcgaagaa catcaagcta ggtgctggtt 3960
tccaagacgt taacgcagaa gataaagtaa gctggaaagg cacgattgac ctgtctaaac 4020
aagacaagcc gtttgacag tctccacaat gtcagacgct aatcacagat ggcacagtca 4080
gtgactttgc tcacctacca agccaactcg cggaagaagt gtogaagatt gttgctgaga 4140
aaggcattcc aaacttccaa gcttggaag atggtttgaa atacagtgc ggcgaaaaag 4200
cgttcgctac agaaaatact cgcgtaaaact tctgggacgt tctgtactgg ggcggtactt 4260
cctcagtgtg cgagtggctc aagaaagggt acgacgtgat tgtttctaac ccagattacg 4320
tgtacatgga tatgccatgc gaagttgacc cgaaagagcg tggttactac tgggcaacac 4380
gtgcaacgga tactcgtaag atgtttggct ttgcaccaga gaacatgcct caaaacgcag 4440
aaacttctgt agatcgcgat ggcaatggct ttactggtaa aggtgaaac gaagcgaaac 4500
ctttctacgg tctatctgca caactttggt ctgagacagt acgtaacgac gagcaatacg 4560
agtacatggt attccctcgc gtcctcgtg ctgctcagcg tgcattggc cgtgctgact 4620
gggaaaacga ctacaaagt ggtgttgagt actcgcaaaa ctctaactta gttgataaag 4680
catcgctaaa ccaagactac aaccgcttg cgaacgtgat tggtaacgt gaactggcta 4740
agctagaaaa atcaggtatt gactaccgcc taccagtacc aggtgcaaaa gtagaagatg 4800
gtaagctagc aatgaacgtt cagttccctg gcgtaacgct tcaatactct ctggatggtg 4860
agaactggtt gacttatgca gacaacgctc gtccaaatgt aactggtgaa gtcttcatcc 4920
gtcgcgtatc tgcgacaggt gagaaggtaa gccgtatcac tagcgtgaaa taatagcgct 4980
cagtattcac taaaatcata gttccttact caaagccctc aacttatgtt gggggctttg 5040
tttatttttc ttcgaaaaat aagcgtgatc atgcgacagg ttgatgaca aaaaattagc 5100
gcaagaagac aaaaatcacc ttgcgctaact gctctgttac aggtcactaa taccatctaa 5160
gtagttgatt catagtgcact gcatatgttg tgttttacag tattatgtag tctgtttttt 5220
atgcaaaatc taatttaata tattgatatt tatatcattt tacgtttctc gttcagcttt 5280
tttatactaa gttggcatta taaaaaagca ttgcttatca atttgttgca acgaacaggt 5340
cactatcagt caaaaataaaa tcattatttg atttcaattt tgtccactc cctgcctctg 5400
tcacacagat actgtgatgc catggtgtcc gacttatgcc cgagaagatg ttgagcaaac 5460
ttatcgctta tctgcttctc atagagtctt gcagacaaac tgcgcaactc gtgaaaggta 5520
ggcgcccccg gtcgaatttg ctttcgaatt tctgccattc atccgcttat tatcacttat 5580
tcaggcgtag caccagggcg ttaagggcac caataactgc cttaaaaaaa ttacgccccg 5640
ccctgccact catcgagta ctgttgtaat tcattaagca ttctgccgac atggaagcca 5700
tcacagacgg catgatgaac ctgaatcgcc agcggcatca gcacctgtc gccttgcgta 5760
taatatttgc ccatggtgaa aacggggcg aagaagttgt ccatattggc cacgtttaaa 5820
tcaaaactgg tgaaactcac ccagggattg gctgagacga aaaacatatt ctcaataaac 5880
cctttaggga aataggccag gttttcaccg taacacgcca catcttgca atatatgtgt 5940
agaaactgcc ggaaactcgtc gtggtattca ctccagagcg atgaaaacgt ttcagtttgc 6000
tcatggaaaa cgggtgaaca aggggtgaaca ctatcccata tcaccagctc accgtctttc 6060
attgccatac g

```

&lt;210&gt; 13

&lt;211&gt; 5641

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; pJMF3

&lt;400&gt; 13

```

gaattccgga tgagcattca tcaggcgggc aagaatgtga ataaaggccg gataaaactt 60
gtgcttattt ttctttacgg tctttaaaaa ggccgtaata tccagctgaa cggctctggtt 120
ataggtaacat tgagcaactg actgaaatgc ctcaaaatgt tctttacgat gccattggga 180
tatataacac gtggtatatac cagtattttt tttctccatt ttagcttcct tagctcctga 240
aaatctcgat aactcaaaaa atacgccccg tagtgatctt atttcattat ggtgaaagtt 300
ggaacctctt acgtgccgat caacgtctca ttttcgcaa aagttggccc agggcttccc 360
ggtatcaaca gggacaccag gatttattta ttctgcgaag tgatcttcg tcacaggtat 420
ttattcggcg caaagtgcgt cgggtgatgc tgccaactta ctgatttagt gtatgatggt 480
gtttttgagg tgctocagtg gcttctgttt ctatcagctg tccctcctgt tcagctactg 540
acggggtggt gcgtaacggc aaaagcaccg ccggacatca gcgtagcgg agtgtgcggc 600
cgactggct tactatgttg gactgatga ggggtcagt gaagtgttc atgtggcagg 660
agaaaaaagg ctgcaccggt gcgtcagcag aatatgtgat acaggatata ttccgcttcc 720

```

tcgctcactg	actcgctacg	ctcggtcggt	cgactgcggc	gagcggaaat	ggcttacgaa	780
cggggcgag	atttcctgga	agatgccagg	aagatactta	acagggaaat	gagagggccg	840
cggcaaaagcc	gtttttccat	aggctccgcc	cccctgacaa	gcatacacgaa	atctgacgct	900
caaatcagtg	gtggcgaaac	ccgacaggac	tataaagata	ccaggcggtt	cccctggcgg	960
ctccctcgtg	cgctctcctg	ttcctgcctt	tcggtttacc	ggtgtcattc	cgctgttatg	1020
gccgcgtttg	tctcattcca	cgctgacac	tcagttccgg	gtaggcagtt	cgctccaagc	1080
tggactgtat	gcacgaaccc	cccgttcagt	ccgaccgctg	cgcttatcc	ggtaactatc	1140
gtcttgagtc	caaccgggaa	agacatgcaa	aagcaccact	ggcagcagcc	actggtaatt	1200
gatttagagg	agttagtctt	gaagtcatgc	gccggttaag	gctaaactga	aaggacaagt	1260
tttggtgact	gcgctcctcc	aagccagtta	cctcggttca	aagagttggt	agctcagaga	1320
accttcgaaa	aaccgcctg	caaggcggtt	ttttcgttt	cagagcaaga	gattacgcgc	1380
agacaaaac	gatctcaaga	agatcatctt	atgcggccgc	atcagataaa	atatttctag	1440
atgccgaact	cagaagtga	acgccgtagc	gccgatggta	gtgtggggtc	tccccatgcg	1500
agagtaggga	actgccaggc	atcaaatata	acgaaaggct	cagtcgaaag	actgggcctt	1560
tcgttttatc	tgttgttgt	cggtgaacgc	tctcctgagt	aggacaaatc	cgccgggagc	1620
ggatttgaac	gttgcgaaag	aacggcccgg	agggtggcgg	gcaggacgcc	cgccataaac	1680
tgccaggcat	caaattaaag	agaaggccat	cctgacggat	ggcctttttg	cgtttctaca	1740
aactcttcc	gtcgtcatat	ctacaagcca	tccccgcgat	gcattaatgt	gagttagctc	1800
actcattagg	caccccaggc	tttacacttt	atgcttccgg	ctcgtatgtt	gtgtggaatt	1860
gtgagcggat	aacaatttca	cacaggaaac	agctatgacc	atgattacgc	caagcttgca	1920
tgcctgcagg	tcgactctag	aggatccccg	ggtaccgagc	tctgagcaac	aagttgtaaa	1980
ctcactggct	gataaccttg	atatccaata	tgaagtgtta	actaaccatg	gtgctaacga	2040
aggctctgcg	tgccaagata	tgggcgcaga	atgggcttct	tgtacaaaag	taaacatgac	2100
gcttggtaac	caagtgaaag	ctgttgactc	aaaagattgg	gctatttact	tccacagcat	2160
tcgtctgatt	ctggatgttg	acaacgagca	gttcaaaatc	tctcgtgtaa	cgggtgacct	2220
acataagcta	gaaccaacag	ataagtttga	cggcttcgct	gccgggtgaag	aggttgttct	2280
tccattgggt	ggtgaatact	ggcaactatt	tgaactgac	ttcatgccgg	gtgcattcgt	2340
ttctgctcca	aacgcagaa	ctaagatgat	tgcttctcta	aatactgaag	atgttgctgc	2400
ttttgtgacg	ggtcttgaag	gtaacaacct	aaaacgtaca	ccagatgaca	acaatgtatt	2460
tgcaaacgct	gtgtctcgtt	ttgagaaaaa	cgaagacct	gcaacacaag	acgtatcaac	2520
caggttacta	ccaacaccaa	tgcacgttga	agcgggttaa	ggcaaagtag	atatcgcgga	2580
tgttattcgc	ctgcttaaac	acgcattcga	tgcactcag	ttcgacgca	ttcaagatcg	2640
tgcagaagtg	gtagggtggg	acgttcgtgg	tgatcttcc	gtaagcatca	ctgttggtcc	2700
tgcagacttc	accggtgaat	tagcaaaatc	tgggtcttac	gaaatgagca	tcaaaggcga	2760
cggtattgtg	attaaagcgt	tcgaccaagc	aggcgcttcc	tacgcagtac	aatctatctt	2820
tggcctggta	gatagccaaa	atgctgattc	tctaccacaa	ctgtctatta	aagatgcgcc	2880
tcgttttgat	taccgtgggt	tgatgggtga	tgtggctcgt	aacttccact	ctaaggacgc	2940
aatccttgca	aataagcaaa	aaatggcagc	gtacaagatg	aacaaacttc	accttcacct	3000
aaccgatgat	gaaggctggc	gtttagaaat	cccgggtctg	cctgagctga	cagaagtggg	3060
tgctaaccgt	tgtttcgata	cacaagagaa	aagctgttta	ctgcctcagc	ttggctctgg	3120
tccaacgaca	gacaactttg	gctctggcta	cttcagcaaa	gcagactacg	tggaaatctt	3180
gaaatacgcg	aaagcacgta	acattgaagt	gattccagaa	atcgatatgc	cagctcacgc	3240
tcgtgcagca	gtagtatcaa	tggaaagctc	ttacgaccgc	ctaattggaag	aaggtaaaga	3300
agctgaagcg	aacgaatacc	gtctgatgga	tcctcaagat	acatcaaacg	taacgacggt	3360
tcagttctac	aataagcaaa	gcttcatcaa	ccatgtatg	gaatcttcaa	ctcgtttgt	3420
tgataagggt	atttcagaag	tggcagcaat	gcaccaagaa	gctggcgctc	cactaacaac	3480
ttggcacttc	ggtggtgacg	aagcgaagaa	catcaagcta	ggtgctggtt	tccaagacgt	3540
taacgcagaa	gataaagtaa	gctggaaagg	cacgattgac	ctgtctaaac	aagacaagcc	3600
gtttgcacag	tctccacaat	gtcagacgct	aatcacagat	ggcacagtca	gtgactttgc	3660
tcacctacca	agccacttcg	cggagaaggt	gtcgaagatt	gttgctgaga	aaggcattcc	3720
aaacttccaa	gcttggcaag	atggtttgaa	atacagtgc	ggcgaaaaag	cgttcgtctc	3780
agaaaatact	cgcgtaaact	tctgggacgt	tctgtactgg	ggcggtactt	cctcagtgt	3840
cgagtggtct	aagaaagggt	acgacgtgat	tgtttctaac	ccagattacg	tgtacatgga	3900
tatgccatac	gaagttgacc	cgaaagagcg	tggttactac	tgggcaaac	gtgcaacgga	3960
tactcgtaag	atgtttggct	ttgcaccaga	gaacatgcct	caaaacgcag	aaacttctgt	4020
agatcgcgat	ggcaatggct	ttactggtaa	aggtgaaatc	gaagcgaaac	ctttctacgg	4080
tctatctgca	caactttggt	ctgagacagt	acgtaacgac	gagcaatacg	agtacatggt	4140
attccctcgc	gtcctcgtcg	ctgctcagcg	tgcattggca	cgtgctgact	gggaaaaacga	4200
ctcaagaagt	ggtggtgagt	actcgcaaaa	ctctaactca	gttgataaag	catcgctaaa	4260
ccaagactac	acgcgctttg	cgaacgact	tgtcaacgct	gaactggcta	agctagaaaa	4320
atcagggtatt	gactaccgcc	taccagtacc	aggtgcaaaa	gtagaagatg	gtaagctagc	4380

aatgaacggt	cagttccctg	gcgtaacgct	tcaatactct	ctggatggtg	agaactgggt	4440
gacttatgca	gacaacgctc	gtccaaatgt	aactgggtgaa	gtcttcatcc	gctcgggtatc	4500
tgcgacaggt	gagaaggtaa	gccgtatcac	tagcgtgaaa	taatagcgct	cagtattcac	4560
taaaatcata	gttccttact	caaagccctc	aacttatggt	gggggctttg	ttatttttcc	4620
ttcggaaaat	aagcgtgatc	atgcgacagg	tttgatgaca	aaaaattagc	gcaagaagac	4680
aaaaatcacc	ttgcgctaatt	gctctgttac	aggctactaa	taccatctaa	gtagttgatt	4740
catagtgact	gcatatgttg	tgttttacag	tattatgtag	tctgtttttt	atgcaaaatc	4800
taatttaata	tattgatatt	tatatcattt	tacgtttctc	gttcagcttt	tttatactaa	4860
gttggcatta	taaaaaagca	ttgcttatca	atttggtgca	acgaacagg	cactatcagt	4920
caaaataaaa	tcattatttg	atttcaattt	tgtccactc	cctgcctctg	tcatacagat	4980
actgtgatgc	catggtgtcc	gacttatgcc	cgagaagatg	ttgagcaaac	ttatcgctta	5040
tctgcttctc	atagagtctt	gcagacaaac	tgcgcaactc	gtgaaaggta	ggcgcccccg	5100
gtcgaatttg	ctttcgaatt	tctgccattc	atccgcttat	tatcacttat	tcaggcgtag	5160
caccaggcgt	ttaagggcac	caataactgc	cttaaaaaaa	ttacgccccg	ccctgccact	5220
catcgtagta	ctgttgtaatt	tcattaagca	ttctgcccag	atggaagcca	tcacagacgg	5280
catgatgaac	ctgaatcgcc	agcggcatca	gcacctgtgc	gccttgcgta	taatatattgc	5340
ccatggtgaa	aacggggggc	aagaagttgt	ccatattggc	cacgtttaaa	tcaaaactgg	5400
tgaaactcac	ccagggtattg	gctgagacga	aaaacatatt	ctcaataaac	cctttaggga	5460
aataggccag	gttttcaccg	taacacgcc	catcttgcca	atatatgtgt	agaaactgcc	5520
ggaaatcgtc	gtggtattca	ctccagagcg	atgaaaacgt	ttcagtttgc	tcattggaaaa	5580
cggtgtaaca	aggggtgaaca	ctatcccata	tcaccagctc	accgtctttc	attgccatac	5640
g						5641

&lt;210&gt; 14

&lt;211&gt; 5670

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; pJMF4

&lt;400&gt; 14

gaattccgga	tgagcattca	tcaggcgggc	aagaatgtga	ataaaggccg	gataaaactt	60
gtgcttattt	ttctttacgg	tctttaaaaa	ggcogtaata	tccagctgaa	cggtctggtt	120
ataggtacat	tgagcaactg	actgaaatgc	ctcaaaatgt	tctttacgat	gccattggga	180
tatatcaacg	gtggtatatt	cagtgatatt	tttctccatt	ttagcttcct	tagctcctga	240
aaatctcgat	aactcaaaaa	atacgcccgg	tagtgatctt	atttcattat	ggtgaaagtt	300
ggaacctctt	acgtgccgat	caacgtctca	ttttcgccaa	aagttggccc	agggtctccc	360
ggtatcaaca	gggacaccag	gatttattta	ttctgcgaag	tgatcttccg	tcacagggtat	420
ttattcggcg	caaagtgcgt	cgggtgatgc	tgccaaactta	ctgatttagt	gtatgatggt	480
gtttttgagg	tgctccagtg	gcttctgttt	ctatcagctg	tccctcctgt	tcagctactg	540
acggggtggt	gcgtaacggc	aaaagcaccg	ccggacatca	gcgctagcgg	agtgtgcggc	600
cgcactggct	tactatgttg	gcactgatga	gggtgtcagt	gaagtgtctc	atgtggcagg	660
agaaaaaagg	ctgcaccggg	gcgtcagcag	aatatgtgat	acaggatata	ttccgcttcc	720
tcgctcactg	actcgctacg	ctcggtcggt	cgactgcggc	gagcggaaat	ggcttacgaa	780
cggggcgagg	atttcctgga	agatgccagg	aagatactta	acagggaagt	gagagggccg	840
cggcaaaagg	gtttttccat	aggctccgcc	cccctgacaa	gcatacagaa	atctgacgct	900
caaatcagtg	gtggcgaaac	ccgacaggac	tataaagata	ccaggcggtt	cccctggcgg	960
ctccctcgtg	cgctctcctg	ttcctgcctt	tcggtttacc	ggtgtcattc	cgctgttatg	1020
gccgcgtttg	tctcattcca	cgcttgacac	tcagttccgg	gtaggcagtt	cgctccaagc	1080
tggaactgtat	gcacgaaccc	cccgttcagt	ccgaccgctg	cgcttatccc	ggtaactatc	1140
gtcttgagtc	caaccgcgaa	agacatgcaa	aagcaccact	ggcagcagcc	actggttaatt	1200
gatttagagg	agttagtctt	gaagtcatgc	gccggttaag	gctaaactga	aaggacaagt	1260
tttggtgact	gcgtccctcc	aagccagtta	cctcggttca	aagagttggt	agctcagaga	1320
accttcgaaa	aaccgcctcg	caaggcgggt	ttttcggttt	cagagcaaga	gattacgcgc	1380
agaccaaaaa	gatctcaaga	agatcatctt	atgcggccgc	atcagataaa	atatttctag	1440
atgccgaact	cagaagtga	acgcctgtag	gccggtggtg	gtgtggggtc	tccccatgcg	1500
agagttagga	actgccaggc	atcaaaataa	acgaaaggct	cagtcgaaag	actgggcctt	1560
tcgttttatc	tggtgtttgt	cgggtgaacg	tctcctgagt	aggacaaatc	cgccgggagc	1620
ggatttgaa	gttgcgaaag	aacggcccg	aggggtggcg	gcaggacgcc	cgccataaac	1680
tgccaggcat	caaattaagg	agaaggccat	cctgacggat	ggcctttttg	cgtttctaca	1740

aactcttctc	gtcgtcatat	ctacaagcca	tcccccgcat	gcattaatgt	gagttagctc	1800
actcattagg	caccccaggc	tttacacttt	atgcttccgg	ctcgtatgtt	gtgtggaatt	1860
gtgagcggat	aacaatttca	cacaggaaac	agctatgacc	atgattacgc	caagcttgca	1920
tgctgcagg	tcgactctag	aggatccccg	ggtaccgagc	tctgagcaac	aagttgtaaa	1980
ctcactggct	gataaccttg	atatccaata	tgaagtgtta	actaaccatg	gtgctaacga	2040
aggtcttgcg	tgccaagata	tgggcgcaga	atgggcttct	tgtatacaaa	taaacatgac	2100
gcttgtaaac	caaggtgaag	ctggtgactc	aaaagattgg	gctattttact	tccacagcat	2160
tcgtctgatt	ctggatgttg	acaacgagca	gttcaaaatc	tctcgtgtaa	cgggtgacat	2220
acataagcta	gaaccaacag	ataagtttga	cggcttcgct	gccggtgaag	agggtgttct	2280
tccattgggt	gggtgaatac	ggcaactatt	tgaactgac	ttcatgccgg	gtgcattcgt	2340
ttctgtccca	aacgcagaac	ctaagatgat	tgcttctcta	aatactgaag	atgttgcgctc	2400
ttttgtgacg	ggtcttgaa	gtaacaacct	aaaacgtaca	ccagatgaca	acaatgtatt	2460
tgcaaacgct	gtgtctcgtt	ttgagaaaaa	cgaagaccta	gcaacacaag	acgtatcaac	2520
cacgttacta	ccaacaccaa	tgacggttga	agcgggtaaa	ggcaaaagtag	atatcgcgga	2580
tggtattgcg	ctgcctaaag	acgcattcga	tgcgactcag	ttcgcagcga	ttcaagatcg	2640
tgcaagaagt	gtagggtg	acgttcgtgg	tgatcttctc	gtaagcatca	ctgttggtcc	2700
tgcaacttc	accggtgaat	tagcaaaatc	tgggtgcttac	gaaatgagca	tcaaaggcga	2760
cggatttg	attaaagcgt	tcgaccaagc	aggcgcttct	tacgcagtag	aatctatctt	2820
tgccctggtg	gatagccaaa	atgctgattc	tctaccacaa	ctgtctatta	aagatgcgcc	2880
tcgttttgat	taccgtggtg	tgatgggtga	tgtggctcgt	aacttccact	ctaaggacgc	2940
aatccttgca	acgctagacc	aaatggcagc	gtacaagatg	aacaaacttc	accttcacct	3000
aaccgatgat	gaaggctggc	gtttagaaat	cccggtctcg	cctgagctga	cagaagtggg	3060
tgctaaccgt	tgtttcgata	cacaagagaa	aagctgttta	ctgcctcagc	ttggctctgg	3120
tccaacgaca	gacaactttg	gctctggcta	cttcagcaaa	gcagactacg	tggaaatctt	3180
gaaatacgcg	aaagcacgta	acattgaagt	gattccagaa	atcgatatgc	cagctcacgc	3240
tcgtgcagca	gtagtatcaa	tggaaagctcg	ttacgaccgc	ctaataggaag	aaggtaaaaga	3300
agctgaagcg	aacgaatacc	gtctgatgga	tcctcaagat	acatcaaacg	taacgacggt	3360
tcagttctac	aataagcaaa	gcttcatcaa	cccatgtatg	gaatcttcaa	ctcgttttgt	3420
tgataaggtg	atttcagaag	tggcagcaat	gcaccaagaa	gctggcgctc	cactaacaac	3480
ttggcaacttc	gggtggtgacg	aagcgaagaa	catcaagcta	gggtgctggtt	tccaagacgt	3540
taacgcagaa	gataaagtaa	gctggaaagg	cacgattgac	ctgtctaaac	aagacaagcc	3600
gtttgcacag	tctccacaat	gtcagacgct	aatcacagat	ggcacagtcg	gtgactttgc	3660
tcacctacca	agccacttcg	cggagaagat	gtcgaagatt	gttgctgaga	aaggcattcc	3720
aaacttccaa	gcttggaag	atggtttgaa	atacagtgac	ggcgaaaaag	cgttcgctac	3780
agaaaaatac	cgcgtaaact	tctgggacgt	tctgtactgg	ggcggtactt	cctcagtgtg	3840
cgagtggctc	aagaaaggtt	acgacgtgat	tgtttctaac	ccagattacg	tgtacatgga	3900
tatgccatac	gaagttgacc	cgaagagcgc	tggttactac	tgggcaacac	gtgcaacgga	3960
tactcgtaag	atgtttggct	ttgcaccaga	gaacatgcct	caaaacgcag	aaacttctgt	4020
agatcgcgat	ggcaatggct	ttactggtaa	aggtgaaatc	gaagcgaaac	ctttctacgg	4080
tctatctgca	caactttggt	ctgagacagt	acgtaacgac	gagcaatagc	agtacatggt	4140
attccctcgc	gtcctcgtcg	ctgctcagcg	tgcattggc	cgtgctgact	gggaaaacga	4200
ctacaaagtt	ggtgttgagt	actcgcaaaa	ctctaactca	gttgataaag	catcgctaaa	4260
ccaagactac	aaccgcttgc	cgaacgtact	tggccaacgt	gaactggcta	agctagaaaa	4320
atcaggtatt	gactaccgcc	taccagtacc	aggtgcaaaa	gtagaagatg	gtaagctagc	4380
aatgaacggt	cagttccctg	gcgtaacgct	tcaatactct	ctggatgggtg	agaactgggt	4440
gacttatgca	gacaacgctc	gtccaaatgt	aactgggtgaa	gtcttcatcc	gctcgggtatc	4500
tgcgacaggt	gagaaggtaa	gccgtatcac	tagcgtgaaa	taatagcgct	cagtattcac	4560
taaaatcata	gttccttact	caaagccctc	aacttatgtt	gggggctttg	tttatttttc	4620
ttcggaaaaa	aagcgtgac	ccccgggggc	gcctaccttt	cacgagttgc	gcagtttgct	4680
tgcaagactc	tatgagaagc	agataagcga	taagtttgct	caacatcttc	tcgggcataa	4740
gtcggacacc	atggcatcac	agtatcgtga	tgacagagcg	agggagtggg	acaaaattga	4800
aatcaaataa	tgattttatt	ttgactgata	gtgacctgtt	cgttgcaaca	aattgataag	4860
caatgctttt	ttataatgcc	aacttagtat	aaaaaagctg	aacgagaaac	gtaaaatgat	4920
ataaatatca	atatattaaa	ttagattttg	cataaaaaac	agactacata	atactgtaaa	4980
acacaacata	tgacgtcact	atgaatcaac	tacttagatg	gtattagtga	cctgtaacag	5040
agcattagcg	caaggtgatt	tttgtcttct	tgcgctaatt	ttttgtcatc	aaacctgtcg	5100
catatcatg	gggctgcagg	aattcagtag	togaatttgc	tttogaattt	ctgccattca	5160
tccgcttatt	atcacttatt	caggcgtagc	accaggcgtt	taagggcacc	aataactgcc	5220
ttaaaaaaat	tacgccccgc	cctgccactc	atcgacgtac	tgttgtaatt	cattaagcat	5280
tctgccgaca	tggaagccat	cacagacggc	atgatgaacc	tgaatcgcca	gcggcatcag	5340
caccttgcgc	ccttgcgtat	aatatttgc	catggtgaaa	acgggggcga	agaagttgtc	5400

```

catattggcc acgttttaaat caaaactggt gaaactcacc cagggattgg ctgagacgaa 5460
aaacatatattc tcaataaacc ctttagggaa ataggccagg ttttcaccgt aacacgccac 5520
atcttgcgaa tataatgtgta gaaactgccg gaaatcgctg tggattcac tccagagcga 5580
tgaaaacggt tcagtttgct catggaaaac ggtgtaacaa ggtgtaacac tatcccatat 5640
caccagctca ccgtctttca ttgccatacg 5670

```

<210> 15  
 <211> 238  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> LacZ(/chitobiase Fusion

<220>  
 <221> CDS  
 <222> (119)...(240)

<221> -35\_signal  
 <222> (44)...(50)  
 <223> Lac promoter

<221> -10\_signal  
 <222> (69)...(74)  
 <223> Lac promoter  
 <221> protein\_bind  
 <222> (81)...(107)  
 <223> Lac repressor binding site

<221> protein\_bind  
 <222> (7)...(34)  
 <223> CAP-cAMP binding site

```

<400> 15
gcatgcatta atgtgagtta gctcactcat taggcacccc aggcattaca ctttatgctt 60
ccggctcgta tgttggtgg aattgtgagc ggataacaat ttcacacagg aaacagct 118
atg acc atg att acg cca agc ttg cat gcc tgc agg tgc act cta gag 166
Met Thr Met Ile Thr Pro Ser Leu His Ala Cys Arg Ser Thr Leu Glu
1 5 10 15

```

```

gat ccc cgg gta ccg agc tct gag caa caa gtt gta aac tca ctg gct 214
Asp Pro Arg Val Pro Ser Ser Glu Gln Gln Val Val Asn Ser Leu Ala
20 25 30

```

```

gat aac ctt gat atc caa tat gaa 238
Asp Asn Leu Asp Ile Gln Tyr Glu
35 40

```

<210> 16  
 <211> 40  
 <212> PRT  
 <213> Artificial Sequence  
 <220>  
 <223> LacZ(/chitobiase Fusion

```

<400> 16
Met Thr Met Ile Thr Pro Ser Leu His Ala Cys Arg Ser Thr Leu Glu
1 5 10 15
Asp Pro Arg Val Pro Ser Ser Glu Gln Gln Val Val Asn Ser Leu Ala
20 25 30
Asp Asn Leu Asp Ile Gln Tyr Glu

```

35

40

&lt;210&gt; 17

&lt;211&gt; 22

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; dnaA/chitobiase Fusion

&lt;400&gt; 17

Met Ser Leu Ser Leu Trp Gln Gln Cys Leu Ala Arg Leu Gln Asp Glu

1

5

10

15

Leu Val Pro Ser Ser Glu

20

&lt;210&gt; 18

&lt;211&gt; 2643

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)...(2643)

&lt;223&gt; dnaA/Chitobiase Fusion

&lt;400&gt; 18

gtg tca ctt tcg ctt tgg cag cag tgt ctt gcc cga ttg cag gat gag 48

Met Ser Leu Ser Leu Trp Gln Gln Cys Leu Ala Arg Leu Gln Asp Glu

1

5

10

15

ttg gta ccg agc tct gag caa caa gtt gta aac tca ctg gct gat aac 96

Leu Val Pro Ser Ser Glu Gln Gln Val Val Asn Ser Leu Ala Asp Asn

20

25

30

ctt gat atc caa tat gaa gtg tta act aac cat ggt gct aac gaa ggt 144

Leu Asp Ile Gln Tyr Glu Val Leu Thr Asn His Gly Ala Asn Glu Gly

35

40

45

ctt gcg tgc caa gat atg ggc gca gaa tgg gct tct tgt aac aaa gta 192

Leu Ala Cys Gln Asp Met Gly Ala Glu Trp Ala Ser Cys Asn Lys Val

50

55

60

aac atg acg ctt gtt aac caa ggt gaa gct gtt gac tca aaa gat tgg 240

Asn Met Thr Leu Val Asn Gln Gly Glu Ala Val Asp Ser Lys Asp Trp

65

70

75

80

gct att tac ttc cac agc att cgt ctg att ctg gat gtt gac aac gag 288

Ala Ile Tyr Phe His Ser Ile Arg Leu Ile Leu Asp Val Asp Asn Glu

85

90

95

cag ttc aaa atc tct cgt gta acg ggt gac cta cat aag cta gaa cca 336

Gln Phe Lys Ile Ser Arg Val Thr Gly Asp Leu His Lys Leu Glu Pro

100

105

110

aca gat aag ttt gac ggc ttc gct gcc ggt gaa gag gtt gtt ctt cca 384

Thr Asp Lys Phe Asp Gly Phe Ala Ala Gly Glu Glu Val Val Leu Pro

115

120

125

ttg gtt ggt gaa tac tgg caa cta ttt gaa act gac ttc atg ccg ggt 432

Leu Val Gly Glu Tyr Trp Gln Leu Phe Glu Thr Asp Phe Met Pro Gly

130	135	140	
gca ttc gtt tct gct cca aac gca gaa cct aag atg att gct tct cta Ala Phe Val Ser Ala Pro Asn Ala Glu Pro Lys Met Ile Ala Ser Leu 145 150 155 160			480
aat act gaa gat gtt gcg tct ttt gtg acg ggt ctt gaa ggt aac aac Asn Thr Glu Asp Val Ala Ser Phe Val Thr Gly Leu Glu Gly Asn Asn 165 170 175			528
cta aaa cgt aca cca gat gac aac aat gta ttt gca aac gct gtg tct Leu Lys Arg Thr Pro Asp Asp Asn Asn Val Phe Ala Asn Ala Val Ser 180 185 190			576
cgt ttt gag aaa aac gaa gac cta gca aca caa gac gta tca acc acg Arg Phe Glu Lys Asn Glu Asp Leu Ala Thr Gln Asp Val Ser Thr Thr 195 200 205			624
tta cta cca aca cca atg cac gtt gaa gcg ggt aaa ggc aaa gta gat Leu Leu Pro Thr Pro Met His Val Glu Ala Gly Lys Gly Lys Val Asp 210 215 220			672
atc gcg gat ggt att gcg ctg cct aaa gac gca ttc gat gcg act cag Ile Ala Asp Gly Ile Ala Leu Pro Lys Asp Ala Phe Asp Ala Thr Gln 225 230 235 240			720
ttc gca gcg att caa gat cgt gca gaa gtg gta ggt gtg gac gtt cgt Phe Ala Ala Ile Gln Asp Arg Ala Glu Val Val Gly Val Asp Val Arg 245 250 255			768
ggt gat ctt cct gta agc atc act gtt gtt cct gca gac ttc acc ggt Gly Asp Leu Pro Val Ser Ile Thr Val Val Pro Ala Asp Phe Thr Gly 260 265 270			816
gaa tta gca aaa tct ggt gct tac gaa atg agc atc aaa ggc gac ggt Glu Leu Ala Lys Ser Gly Ala Tyr Glu Met Ser Ile Lys Gly Asp Gly 275 280 285			864
att gtg att aaa gcg ttc gac caa gca ggc gct ttc tac gca gta caa Ile Val Ile Lys Ala Phe Asp Gln Ala Gly Ala Phe Tyr Ala Val Gln 290 295 300			912
tct atc ttt ggc ctg gta gat agc caa aat gct gat tct cta cca caa Ser Ile Phe Gly Leu Val Asp Ser Gln Asn Ala Asp Ser Leu Pro Gln 305 310 315 320			960
ctg tct att aaa gat gcg cct cgt ttt gat tac cgt ggt gtg atg gtg Leu Ser Ile Lys Asp Ala Pro Arg Phe Asp Tyr Arg Gly Val Met Val 325 330 335			1008
gat gtg gct cgt aac ttc cac tct aag gac gca atc ctt gca acg cta Asp Val Ala Arg Asn Phe His Ser Lys Asp Ala Ile Leu Ala Thr Leu 340 345 350			1056
gac caa atg gca gcg tac aag atg aac aaa ctt cac ctt cac cta acc Asp Gln Met Ala Ala Tyr Lys Met Asn Lys Leu His Leu His Leu Thr 355 360 365			1104
gat gat gaa ggc tgg cgt tta gaa atc ccg ggt ctg cct gag ctg aca Asp Asp Glu Gly Trp Arg Leu Glu Ile Pro Gly Leu Pro Glu Leu Thr 370 375 380			1152

gaa gtg ggt gct aac cgt tgt ttc gat aca caa gag aaa agc tgt tta	1200
Glu Val Gly Ala Asn Arg Cys Phe Asp Thr Gln Glu Lys Ser Cys Leu	
385 390 395 400	
ctg cct cag ctt ggc tct ggt cca acg aca gac aac ttt ggc tct ggc	1248
Leu Pro Gln Leu Gly Ser Gly Pro Thr Thr Asp Asn Phe Gly Ser Gly	
405 410 415	
tac ttc agc aaa gca gac tac gtg gaa atc ttg aaa tac gcg aaa gca	1296
Tyr Phe Ser Lys Ala Asp Tyr Val Glu Ile Leu Lys Tyr Ala Lys Ala	
420 425 430	
cgt aac att gaa gtg att cca gaa atc gat atg cca gct cac gct cgt	1344
Arg Asn Ile Glu Val Ile Pro Glu Ile Asp Met Pro Ala His Ala Arg	
435 440 445	
gca gca gta gta tca atg gaa gct cgt tac gac cgc cta atg gaa gaa	1392
Ala Ala Val Val Ser Met Glu Ala Arg Tyr Asp Arg Leu Met Glu Glu	
450 455 460	
ggg aaa gaa gct gaa gcg aac gaa tac cgt ctg atg gat cct caa gat	1440
Gly Lys Glu Ala Glu Ala Asn Glu Tyr Arg Leu Met Asp Pro Gln Asp	
465 470 475 480	
aca tca aac gta acg acg gtt cag ttc tac aat aag caa agc ttc atc	1488
Thr Ser Asn Val Thr Thr Val Gln Phe Tyr Asn Lys Gln Ser Phe Ile	
485 490 495	
aac cca tgt atg gaa tct tca act cgc ttt gtt gat aag gtg att tca	1536
Asn Pro Cys Met Glu Ser Ser Thr Arg Phe Val Asp Lys Val Ile Ser	
500 505 510	
gaa gtg gca gca atg cac caa gaa gct ggc gct cca cta aca act tgg	1584
Glu Val Ala Ala Met His Gln Glu Ala Gly Ala Pro Leu Thr Thr Trp	
515 520 525	
cac ttc ggt ggt gac gaa gcg aag aac atc aag cta ggt gct ggt ttc	1632
His Phe Gly Gly Asp Glu Ala Lys Asn Ile Lys Leu Gly Ala Gly Phe	
530 535 540	
caa gac gtt aac gca gaa gat aaa gta agc tgg aaa ggc acg att gac	1680
Gln Asp Val Asn Ala Glu Asp Lys Val Ser Trp Lys Gly Thr Ile Asp	
545 550 555 560	
ctg tct aaa caa gac aag ccg ttt gca cag tct cca caa tgt cag acg	1728
Leu Ser Lys Gln Asp Lys Pro Phe Ala Gln Ser Pro Gln Cys Gln Thr	
565 570 575	
cta atc aca gat ggc aca gtc agt gac ttt gct cac cta cca agc cac	1776
Leu Ile Thr Asp Gly Thr Val Ser Asp Phe Ala His Leu Pro Ser His	
580 585 590	
ttc gcg gaa gaa gtg tcg aag att gtt gct gag aaa ggc att cca aac	1824
Phe Ala Glu Glu Val Ser Lys Ile Val Ala Glu Lys Gly Ile Pro Asn	
595 600 605	
ttc caa gct tgg caa gat ggt ttg aaa tac agt gac ggc gaa aaa gcg	1872
Phe Gln Ala Trp Gln Asp Gly Leu Lys Tyr Ser Asp Gly Glu Lys Ala	
610 615 620	



ttc gct aca gaa aat act cgc gta aac ttc tgg gac gtt ctg tac tgg	1920
Phe Ala Thr Glu Asn Thr Arg Val Asn Phe Trp Asp Val Leu Tyr Trp	
625 630 635 640	
ggc ggt act tcc tca gtg tac gag tgg tct aag aaa ggt tac gac gtg	1968
Gly Gly Thr Ser Ser Val Tyr Glu Trp Ser Lys Lys Gly Tyr Asp Val	
645 650 655	
att gtt tct aac cca gat tac gtg tac atg gat atg cca tac gaa gtt	2016
Ile Val Ser Asn Pro Asp Tyr Val Tyr Met Asp Met Pro Tyr Glu Val	
660 665 670	
gac ccg aaa gag cgt ggt tac tac tgg gca aca cgt gca acg gat act	2064
Asp Pro Lys Glu Arg Gly Tyr Trp Ala Thr Arg Ala Thr Asp Thr	
675 680 685	
cgt aag atg ttt ggc ttt gca cca gag aac atg cct caa aac gca gaa	2112
Arg Lys Met Phe Gly Phe Ala Pro Glu Asn Met Pro Gln Asn Ala Glu	
690 695 700	
act tct gta gat cgc gat ggc aat ggc ttt act ggt aaa ggt gaa atc	2160
Thr Ser Val Asp Arg Asp Gly Asn Gly Phe Thr Gly Lys Gly Glu Ile	
705 710 715 720	
gaa gcg aaa cct ttc tac ggt cta tct gca caa ctt tgg tct gag aca	2208
Glu Ala Lys Pro Phe Tyr Gly Leu Ser Ala Gln Leu Trp Ser Glu Thr	
725 730 735	
gta cgt aac gac gag caa tac gag tac atg gta ttc cct cgc gtc ctc	2256
Val Arg Asn Asp Glu Gln Tyr Glu Tyr Met Val Phe Pro Arg Val Leu	
740 745 750	
gct gct gct cag cgt gca tgg cac cgt gct gac tgg gaa aac gac tac	2304
Ala Ala Ala Gln Arg Ala Trp His Arg Ala Asp Trp Glu Asn Asp Tyr	
755 760 765	
aaa gtt ggt gtt gag tac tcg caa aac tct aat cta gtt gat aaa gca	2352
Lys Val Gly Val Glu Tyr Ser Gln Asn Ser Asn Leu Val Asp Lys Ala	
770 775 780	
tcg cta aac caa gac tac aac cgc ttt gcg aac gta ctt ggt caa cgt	2400
Ser Leu Asn Gln Asp Tyr Asn Arg Phe Ala Asn Val Leu Gly Gln Arg	
785 790 795 800	
gaa ctg gct aag cta gaa aaa tca ggt att gac tac cgc cta cca gta	2448
Glu Leu Ala Lys Leu Glu Lys Ser Gly Ile Asp Tyr Arg Leu Pro Val	
805 810 815	
cca ggt gca aaa gta gaa gat ggt aag cta gca atg aac gtt cag ttc	2496
Pro Gly Ala Lys Val Glu Asp Gly Lys Leu Ala Met Asn Val Gln Phe	
820 825 830	
cct ggc gta acg ctt caa tac tct ctg gat ggt gag aac tgg ttg act	2544
Pro Gly Val Thr Leu Gln Tyr Ser Leu Asp Gly Glu Asn Trp Leu Thr	
835 840 845	
tat gca gac aac gct cgt cca aat gta act ggt gaa gtc ttc atc cgc	2592
Tyr Ala Asp Asn Ala Arg Pro Asn Val Thr Gly Glu Val Phe Ile Arg	
850 855 860	
tcg gta tct gcg aca ggt gag aag gta agc cgt atc act agc gtg aaa	2640

Ser Val Ser Ala Thr Gly Glu Lys Val Ser Arg Ile Thr Ser Val Lys  
 865 870 875 880

taa  
 \*

2643

<210> 19  
 <211> 880  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> dnaA/Chitobiase Fusion

<400> 19  
 Met Ser Leu Ser Leu Trp Gln Gln Cys Leu Ala Arg Leu Gln Asp Glu  
 1 5 10 15  
 Leu Val Pro Ser Ser Glu Gln Gln Val Val Asn Ser Leu Ala Asp Asn  
 20 25 30  
 Leu Asp Ile Gln Tyr Glu Val Leu Thr Asn His Gly Ala Asn Glu Gly  
 35 40 45  
 Leu Ala Cys Gln Asp Met Gly Ala Glu Trp Ala Ser Cys Asn Lys Val  
 50 55 60  
 Asn Met Thr Leu Val Asn Gln Gly Glu Ala Val Asp Ser Lys Asp Trp  
 65 70 75 80  
 Ala Ile Tyr Phe His Ser Ile Arg Leu Ile Leu Asp Val Asp Asn Glu  
 85 90 95  
 Gln Phe Lys Ile Ser Arg Val Thr Gly Asp Leu His Lys Leu Glu Pro  
 100 105 110  
 Thr Asp Lys Phe Asp Gly Phe Ala Ala Gly Glu Glu Val Val Leu Pro  
 115 120 125  
 Leu Val Gly Glu Tyr Trp Gln Leu Phe Glu Thr Asp Phe Met Pro Gly  
 130 135 140  
 Ala Phe Val Ser Ala Pro Asn Ala Glu Pro Lys Met Ile Ala Ser Leu  
 145 150 155 160  
 Asn Thr Glu Asp Val Ala Ser Phe Val Thr Gly Leu Glu Gly Asn Asn  
 165 170 175  
 Leu Lys Arg Thr Pro Asp Asp Asn Asn Val Phe Ala Asn Ala Val Ser  
 180 185 190  
 Arg Phe Glu Lys Asn Glu Asp Leu Ala Thr Gln Asp Val Ser Thr Thr  
 195 200 205  
 Leu Leu Pro Thr Pro Met His Val Glu Ala Gly Lys Gly Lys Val Asp  
 210 215 220  
 Ile Ala Asp Gly Ile Ala Leu Pro Lys Asp Ala Phe Asp Ala Thr Gln  
 225 230 235 240  
 Phe Ala Ala Ile Gln Asp Arg Ala Glu Val Val Gly Val Asp Val Arg  
 245 250 255  
 Gly Asp Leu Pro Val Ser Ile Thr Val Val Pro Ala Asp Phe Thr Gly  
 260 265 270  
 Glu Leu Ala Lys Ser Gly Ala Tyr Glu Met Ser Ile Lys Gly Asp Gly  
 275 280 285  
 Ile Val Ile Lys Ala Phe Asp Gln Ala Gly Ala Phe Tyr Ala Val Gln  
 290 295 300  
 Ser Ile Phe Gly Leu Val Asp Ser Gln Asn Ala Asp Ser Leu Pro Gln  
 305 310 315 320  
 Leu Ser Ile Lys Asp Ala Pro Arg Phe Asp Tyr Arg Gly Val Met Val  
 325 330 335  
 Asp Val Ala Arg Asn Phe His Ser Lys Asp Ala Ile Leu Ala Thr Leu  
 340 345 350  
 Asp Gln Met Ala Ala Tyr Lys Met Asn Lys Leu His Leu His Leu Thr  
 355 360 365

Asp Asp Glu Gly Trp Arg Leu Glu Ile Pro Gly Leu Pro Glu Leu Thr  
 370 375 380  
 Glu Val Gly Ala Asn Arg Cys Phe Asp Thr Gln Glu Lys Ser Cys Leu  
 385 390 395 400  
 Leu Pro Gln Leu Gly Ser Gly Pro Thr Thr Asp Asn Phe Gly Ser Gly  
 405 410 415  
 Tyr Phe Ser Lys Ala Asp Tyr Val Glu Ile Leu Lys Tyr Ala Lys Ala  
 420 425 430  
 Arg Asn Ile Glu Val Ile Pro Glu Ile Asp Met Pro Ala His Ala Arg  
 435 440 445  
 Ala Ala Val Val Ser Met Glu Ala Arg Tyr Asp Arg Leu Met Glu Glu  
 450 455 460  
 Gly Lys Glu Ala Glu Ala Asn Glu Tyr Arg Leu Met Asp Pro Gln Asp  
 465 470 475 480  
 Thr Ser Asn Val Thr Thr Val Gln Phe Tyr Asn Lys Gln Ser Phe Ile  
 485 490 495  
 Asn Pro Cys Met Glu Ser Ser Thr Arg Phe Val Asp Lys Val Ile Ser  
 500 505 510  
 Glu Val Ala Ala Met His Gln Glu Ala Gly Ala Pro Leu Thr Thr Trp  
 515 520 525  
 His Phe Gly Gly Asp Glu Ala Lys Asn Ile Lys Leu Gly Ala Gly Phe  
 530 535 540  
 Gln Asp Val Asn Ala Glu Asp Lys Val Ser Trp Lys Gly Thr Ile Asp  
 545 550 555 560  
 Leu Ser Lys Gln Asp Lys Pro Phe Ala Gln Ser Pro Gln Cys Gln Thr  
 565 570 575  
 Leu Ile Thr Asp Gly Thr Val Ser Asp Phe Ala His Leu Pro Ser His  
 580 585 590  
 Phe Ala Glu Glu Val Ser Lys Ile Val Ala Glu Lys Gly Ile Pro Asn  
 595 600 605  
 Phe Gln Ala Trp Gln Asp Gly Leu Lys Tyr Ser Asp Gly Glu Lys Ala  
 610 615 620  
 Phe Ala Thr Glu Asn Thr Arg Val Asn Phe Trp Asp Val Leu Tyr Trp  
 625 630 635 640  
 Gly Gly Thr Ser Ser Val Tyr Glu Trp Ser Lys Lys Gly Tyr Asp Val  
 645 650 655  
 Ile Val Ser Asn Pro Asp Tyr Val Tyr Met Asp Met Pro Tyr Glu Val  
 660 665 670  
 Asp Pro Lys Glu Arg Gly Tyr Tyr Trp Ala Thr Arg Ala Thr Asp Thr  
 675 680 685  
 Arg Lys Met Phe Gly Phe Ala Pro Glu Asn Met Pro Gln Asn Ala Glu  
 690 695 700  
 Thr Ser Val Asp Arg Asp Gly Asn Gly Phe Thr Gly Lys Gly Glu Ile  
 705 710 715 720  
 Glu Ala Lys Pro Phe Tyr Gly Leu Ser Ala Gln Leu Trp Ser Glu Thr  
 725 730 735  
 Val Arg Asn Asp Glu Gln Tyr Glu Tyr Met Val Phe Pro Arg Val Leu  
 740 745 750  
 Ala Ala Ala Gln Arg Ala Trp His Arg Ala Asp Trp Glu Asn Asp Tyr  
 755 760 765  
 Lys Val Gly Val Glu Tyr Ser Gln Asn Ser Asn Leu Val Asp Lys Ala  
 770 775 780  
 Ser Leu Asn Gln Asp Tyr Asn Arg Phe Ala Asn Val Leu Gly Gln Arg  
 785 790 795 800  
 Glu Leu Ala Lys Leu Glu Lys Ser Gly Ile Asp Tyr Arg Leu Pro Val  
 805 810 815  
 Pro Gly Ala Lys Val Glu Asp Gly Lys Leu Ala Met Asn Val Gln Phe  
 820 825 830  
 Pro Gly Val Thr Leu Gln Tyr Ser Leu Asp Gly Glu Asn Trp Leu Thr  
 835 840 845  
 Tyr Ala Asp Asn Ala Arg Pro Asn Val Thr Gly Glu Val Phe Ile Arg

	850					855						860			
Ser	Val	Ser	Ala	Thr	Gly	Glu	Lys	Val	Ser	Arg	Ile	Thr	Ser	Val	Lys
865					870					875					880

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
19 April 2001 (19.04.2001)

PCT

(10) International Publication Number  
**WO 01/27322 A3**(51) International Patent Classification<sup>7</sup>: **C12Q 1/68**(21) International Application Number: **PCT/US00/21048**(22) International Filing Date: **2 August 2000 (02.08.2000)**(25) Filing Language: **English**(26) Publication Language: **English**(30) Priority Data:  
**60/159,221 13 October 1999 (13.10.1999) US**(71) Applicant (for all designated States except US): **ELITRA  
PHARMACEUTICALS, INC. [US/US]; Suite A, 3510  
Dunhill Street, San Diego, CA 92121 (US).**

(72) Inventor; and

(75) Inventor/Applicant (for US only): **ZYSKIND, Judith  
[US/US]; 8514 La Jolla Scenic Drive, La Jolla, CA 92047  
(US).**(74) Agent: **HUNT, Dale, C.; 16th Floor, 620 Newport Center  
Drive, Newport Beach, CA 92660 (US).**(81) Designated States (national): **AE, AG, AL, AM, AT, AT  
(utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA,  
CH, CN, CR, CU, CZ, CZ (utility model), DE, DE (utility  
model), DK, DK (utility model), DM, DZ, EE, EE (utility  
model), ES, FI, FI (utility model), GB, GD, GE, GH, GM,  
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KR (utility  
model), KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG,  
MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD,  
SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT,  
TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.**(84) Designated States (regional): **ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).****Published:**

— with international search report

(88) Date of publication of the international search report:  
**13 December 2001***For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*(54) Title: **CHITOBIASE AS A REPORTER ENZYME**

(57) Abstract: The present invention relates to reporter gene constructs encoding a cytoplasmic form of chitobiase (N,N'-diacetyl-chitobiase) and methods of using these reporter gene constructs. The use of a cytoplasmic form of chitobiase as a reporter enzyme is generally applicable to the study of gene expression in organisms which do not contain N-acetyl-β-D-glucosaminidases.

**WO 01/27322 A3**

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/21048

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12Q C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KALABAT D Y ET AL: "Chitobiase, a new reporter enzyme" BIOTECHNIQUES,US,EATON PUBLISHING, NATICK, vol. 25, no. 6, December 1998 (1998-12), pages 1030-1035, XP002106689 ISSN: 0736-6205 cited in the application the whole document	1-50
A	WO 98 49320 A (AHLUWALIA NAVNEET K ;BENKEL BERNHARD F (CA); FALCONER MARCIA M (CA) 5 November 1998 (1998-11-05) abstract page 1 -page 2 --- -/--	1-50



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

6 July 2001

Date of mailing of the international search report

23/07/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Hart-Davis, J

# INTERNATIONAL SEARCH REPORT

International Application No

Pct/US 00/21048

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DATABASE BIOSIS 'Online!  BIOSCIENCES INFORMATION SERVICE,  PHILADELPHIA, PA, US; March 1998 (1998-03)  PIRUZIAN E S ET AL: "The use of a  thermostable beta-glucanase gene from  Clostridium thermocellum as a reporter  gene in plants."  Database accession no. PREV199800227889  XP002171323  abstract  &amp; MOLECULAR &amp; GENERAL GENETICS,  vol. 257, no. 5, March 1998 (1998-03),  pages 561-567,  ISSN: 0026-8925</p>	1-50

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/21048

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9849320 A	05-11-1998	CA 2203613 A	24-10-1998
		AU 6085698 A	24-11-1998



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**